Index of Authors

A

Abbott, U. K., 96*
Abplanalp, H., 104*, 367
Acar, N., 1166
Acha, M., 1229
Adams, A. W., 47*, 89*, 1348
Adams, C. R., 43*
Adams, M. H., 10*, 13*, 43*
Adams, R. L., 43*
Addison, J. B., 1203
Adefope, N. A., 39*, 82*, 168*
Agar, V., 680
Aguillard, C. D., 1*, 19*
Ahn, D. U., 44*
Aho, P. W., 534, 1804
Akashiba, H., 1625
Akpobome, G. O., 45*
Al-Athari, A. K., 45*, 307, 778
Al-Heeti, H. E., 46*
Allen, A. E., 66*
Allen, P. C., 45*
Alphin, R. L., Jr., 46*, 134*
Al-Soudi, K. A., 46*
Alston-Mills, B., 110*
Alvarez, C., 351
Ameenuddin, A., 1829
Aminul Haque, A.K.M., 94*
Ammerman, C. B., 18*, 41*, 1295, 1585
Ammerman, E., 1*, 46*
Amo, J., 1870
Amyot, E., 1758, 1762
Anderson, K. E., 47*, 1348
Andrews, D. K., 47*
Andrews, G. A., 47*
Andrews, L. D., 2*, 48*, 1414
Andrews, P., 40*, 173*
Ang, C.Y.W., 2*, 112*, 510, 736
Angel, C. R., 1807
Angel, R., 153*
Ansah, G. A., 819
Appelgren, L.-E., 707
Araba, M., 11*
Arbabi, A. R., 1414
Arjona, A. A., 2*, 226
Armstrong, G. E., 126*
Arscott, G. H., 153
Atkinson, J. L., 49*
Augustine, P. C., 49*, 72*, 549
Austic, R. E., 14*, 148*, 172*, 750
Avalos, E., 49*
Avila-Gonzalez, E., 159*

B

Bacon, J. R., 50*, 89*
Bacon, L. D., 50*, 111*, 191, 1525
Bacon, W. L., 51*, 107*, 161*, 166*, 357, 826, 1378, 1388, 1660
Badresingh, V. B., 51*
Bagley, L. G., 956
Bai, Y., 52*
Bailey, C. A., 52*, 88*, 90*, 130*, 1154, 1644
Bailey, J. S., 26*, 52*, 53*, 71*, 920, 928
Bain, S. D., 590
Baker, D. H., 94*, 172*, 232, 341, 652
Baker, R. C., 106*, 138*, 557, 1766
Bakst, M. R., 327, 1461
Ball, H. R., Jr., 53*, 73*, 91*
Ballard, R., 113
Bamberg, H., 1363
Barbour, G. W., 53*
Barbut, S., 1797
Barnhart, H. M., 3*
Bartov, I., 420, 809
Bastien, R. W., 3*, 54*
Batson, D. B., 8*
Baughman, G. R., 12*
Beasley, J., 109*
Beck, M. M., 113*
Becker, R., 1166
Beers, K. W., 3*, 30*
Beitz, D. C., 85*
Bell, D. E., 544
Bengal, I., 420
Benibo, B. S., 54*
Bennett, C. D., 49*
Bennink, M. R., 128*
Benoff, F. H., 4*, 72*, 143*
Berkhoff, H. A., 121*
Bernon, D. E., 1497
Bhanushali, J. K., 16*
Bierlmaier, S. J., 166*
Bietz, D. C., 85*
Bilgili, S. F., 54*, 144*, 513, 845, 897, 1810
Bills, D. F., 168*
Bird, R., 43*
Birrenkott, G. P., 1783
Bitgood, J. J., 102*, 530, 1465
Blackman, S., 88
Blair, M. E., 1281
Blair, R., 1274
Blake, J. P., 175*
Blakewood, E. G., 55*
Blankenship, L. C., 26*, 52*, 71*
Blankenship, M., 10*, 13*
Bloom, S. E., 74*
Blore, P., 40*
Bokanyi, R. P., Jr., 55*, 160*
Bond, P. L., 80*, 163*
Bootwalla, S. M., 55*, 56*, 1218
Bornstein, S., 809
Bottcher, R. W., 8*
Bottje, W. G., 3*, 30*, 171*, 1183
Bower, R. K., 1123
Bowers, J. A., 72
Bowes, V. A., 1102
Bowyer, B. L., 4*
Boyle, M. L., III, 56*
Brachet, P., 56*, 81*
Bradley, F. A., 54*, 886
Braithwaite, L. A., 57*
Brake, J., 5*, 37*, 38*, 57*, 58*, 155*, 163*, 313, 538, 669
Brake, J. T., 38*
Brake, N. P., 669
Branton, S. L., 5*, 25*, 34*, 58*, 64, 73*, 116*, 677, 1239
Breeding, S. W., 5*, 58*
Brewer, C. E., 33*
Bricker, J. M., 33*
Briles, C. O., 6*
Briles, W. E., 111*, 1210
Britton, J. G., 58*, 59*
Britton, W. M., 6*, 18*, 59*, 97*, 98*, 205, 213, 339
Brown, D. R., 47*
Brown, J., 1683
Brown, M. L., 123*
Brown-Borg, H. M., 13*, 59*
Brunström, B., 52
Bryan, T. E., 60*
Buckland, R. B., 819
Buckley, D. J., 1105
Buhr, R. J., 60*, 117*, 173*, 367
Bull, S. A., 1486
Buresh, R. E., 336
Burg, J. G., 1015
Burger, R. E., 158*
Burke, W. H., 117*, 979, 1098
Burton, S. D., 167*
Bush, S., 158*
Bushong, R. D., 143*
Buss, E. G., 1322
Butcher, G. D., 60*, 61*, 174*
Butler, G., 680

C

Cabel, M. C., 6*, 7*, 300, 1725
Calabotta, D. F., 1725
Callender, M. E., 1043, 1050
Calvert, C. C., 61*, 104*, 146*
Camp, L. J., 1058
Campbell, L. D., 61*
Campion, D. R., 64*, 674, 1338
Campo, J. L., 351
Cantor, A. H., 7*, 8*, 19*, 41*, 62*, 137*, 149*, 1157
Capasso, J., 35*
Carawan, R. E., 155*
Carey, J. B., 7*, 62*, 63*
Carlson, C. W., 445
Carpenter, G. H., 102*, 137*
Carsia, R. V., 64*, 105*
Carter, E. R., 63*
Carter, T. A., 8*, 63*
Cartwright, A. L., 64*, 71*, 1338
Cason, J. A., 6*, 64*, 213, 979
Cassens, R. G., 973, 982
Castaldo, D. J., 8*, 24*, 65*, 427
Castanon, F., 65*, 94*, 97*
Caston, L. J., 1236
Catherman, D. R., 8*
Cave, N. A., 78
Cecil, H. C., 327, 1461
Chambers, J. R., 65*, 183, 565, 1497

*Indicates Abstract.

Chang, Y., 155*
Chaplin, S. B., 126
Charles, O. W., 103, 1737
Cheeke, P. R., 153
Chen, B. H., 1154, 1644
Chen, F., 66*
Chen, T. C., 9*, 27*, 112*, 128*
Cheng, K. M., 66*, 127*, 865
Cheng, O.A.T., 109*
Cheng, S., 859, 989
Cheng, S. E., 35*
Cheng, T. K., 66*, 67*
Cherry, J. A., 800
Chin, S. F., 1001
Chiu, S. W., 9*
Chou, H.-F., 470
Christensen, V. L., 9*, 17*, 67*, 92*, 956
Christmas, R. B., 9*, 29*, 67*, 68*, 452
Clanton, C., 66*
Clare, R. A., 68*
Clark, S. R., 116*
Classen, H. L., 68*, 145*
Clower, W. B., 2*, 48*
Cogburn, L. A., 69*, 117*, 168*
Colberg, M., 10*, 13*, 69*
Collison, C. H., 1015
Collisson, E. W., 61*, 69*, 158*
Combs, G. F., Jr., 141*
Conner, S. D., 70*
Cook, M. E., 139*, 174*, 175*, 434, 1717
Cook, R. E., 890
Coon, C. N., 66*, 67*, 78*, 109*, 129*, 135*, 136*, 1306
Cooper, K., 1783
Cooper, R., 17*
Copeland, K.F.T., 70*
Coppings, R., 82*
Corrier, D., 60*
Corrier, D. E., 253
Cotter, P. F., 70*
Coulibaly, M., 76*
Cowen, B. S., 90*, 1250, 1694
Cox, E., 40*, 173*
Cox, N. A., 26*, 52*, 53*, 71*, 921
Craig, J. V., 109*, 848. 1400
Cravener, T. L., 168*
Creger, C. R., 77*, 78*, 253
Crissey, S. D., 1345
Crittenden, L. B., 50*, 191, 1668
Croom, W. J., Jr., 101*
Csallany, A. S., 1814
Cuca-G., M., 159*
Cuchens, M. A., 25*
Cunningham, D. L., 198, 399
Cupo, M. A., 71*
Current, W. L., 720, 730
Curtis, C. E., 10*
Curtis, P., 71*
Curtis, S. E., 1787, 1791

D

Daghir, N., 166*
Dake, R. L., 40*
Dale, N. M., 10*, 11*, 21*, 72*, 1188
Dalrymple, R. H., 73*
Damron, B. L., 11*, 15*, 27*, 1302
Danforth, H. D., 49*, 68*, 72*, 100*, 147*
Danielson, D. M., 163*
Daniloff, J. K., 143*
Darden, J. R., 519, 1111
Davidson, J. N., 70*
Davis, C. E., 11*, 23*
Davis, G. S., 12*, 72*
Davis, V., 855
Dawson, P. L., 73, 715, 1545
Day, E. J., 111*, 130*, 131*
De Groote, G., 950
Dean, W. F., 73*
Deaton, J. W., 5*, 27*, 64, 73*, 375, 378, 677, 1239, 1514
deGraft-Hanson, J. A., 74*
Delany, M. E., 74*
DeLee, J. A., 23*
DeLoach, J. R., 133*
Denbow, D. M., 2*, 20*, 25*, 74*, 114*, 226
Denno, K. M., 75*, 116*
Denton, J. H., 75*, 1269, 1568
Derijcke, J., 950
Devaney, J. A., 549
Dhillon, A. S., 75*
Dhladhla, V., 76*
Diambra, O. H., 76*
Dibner, J. J., 12*, 76*, 1314
Dick, J. W., 26*, 135*
Dickens, J. A., 11*, 12*, 23*, 77*, 1069
Dickson, T. G., 1684
Dieckert, J. W., 77*, 78*, 88*
Dieckert, M. C., 77*, 78*
Dietert, R. R., 74*, 141*
Diez, F. N., 91*
Dingfelder, R. S., 116*
Dixon, J. L., 162*
Doamekpor, S. K., 78*
Doi, Y., 476
Donker, R. A., 79*
Donoghue, D. J., 79*
Douglas, J. H., 80*, 163*
Douglass, L. W., 107*
Doumit, M. E., 121*
Driggers, C. D., 10*, 13*, 80*
Dubbert, W. H., 944
Duke, G. E., 126
Duke, S., 103
Dukes, M. G., 544, 715, 1545
Dunnington, E. A., 20*
Dupuis, L., 81*

E

Earl, L. A., 103*
Eason, P. J., 118*
Eberhart, D. E., 13*, 81*
Echevarria, M. G., 1295, 1585
Edens , F. W., 12*, 13*, 38*, 42*, 59*, 72*, 81*, 174*
Edwards, H. M., Jr., 84*, 113, 150*, 159*, 1436
Egbert, W. R., 54*
Ehmler, T. J., 76*
Ehrich, M., 1229
Eissa, A. I., 84*
Ekhator, N., 82*
El Halawani, M. E., 79*, 86*, 93*, 95*, 127*, 128*, 138*, 1794
Elaghbary, A. M., 82*
Eldeeb, M. A., 82*
Elfar, A. A., 83*
El-Gendy, E. A., 14*, 83*
El-Husseiny, O., 84*
Elissalde, M. H., 99*
Elkin, R. G., 851
Elliot, M. A., 84*
Entrikin, R. K., 367
Escribano, F., 84*, 85*, 153*, 1089
Esteve-Garcia, E., 14*
Etches, R. J., 108*, 117*, 137*, 319
Evangelisti, J. W., 1034

F

Fadly, A. M., 1674
Fagan, W. E., 1105
Fancher, B. I., 1469
Fanguy, R. C., 45*
Farr, A. J., 85*
Farran, M. T., 85*
Fehrer, S. C., 86*, 1794
Fell, R. V., 103
Fenderson, C. L., 39*, 168*
Ferguson, T. M., 3*
Ferket, P. R., 132*, 153*, 585
Fethiere, R., 15*, 86*, 120*
Fetterer, R. H., 111*
Findlay, C., 270
Firman, J. D., 86*, 143*
Fitz-Coy, S. H., 87*, 96*
Fitzgerald, C., 71*
Fletcher, D. L., 25*, 35*, 53*, 117*, 275, 635, 674, 908, 979, 1147, 1549, 1702
Flunker, L. K., 11*, 15*, 1302
Forster, B. W., 1647
Forsythe, R. H., 40*
Foster, D. N., 107*, 699
Fox, M. C., 171*
Frackelton, D. G., 36*
Fredericksen, T. L., 147*
Frobish, R. A., 169*
Frolich, A. A., 502
Froman, D. P., 82*, 103*, 1218
Froning, G. W., 87*, 1536
Froyman, R., 950
Fryer, R. L., 1075
Fuchs, R., 707
Fuller, J., 71*
Fuller, J. C., Jr., 87*

G

Galehouse, D. C., 107*
Galyean, R. D., 162*
Gardiner, E. E., 156, 1020, 1243

Gardner, F. A., 75*, 88*, 91*, 130*, 1269, 1431, 1568
Garlich, J. D., 5*, 15*, 20*
Garrison, K., 88*
Garwood, V. A., 88*
Gast, R. K., 689, 699
Gempesaw, C. M., II, 50*, 89*
George, J. C., 463
Gerken, M., 1363
Gernat, A. G., 89*
Ghazalah, A. A., 84*
Ghodrati, A., 82*
Giambrone, J. J., 89*
Gibson, R. M., 90*
Gilbert, D. L., 35*
Gilbert, J. M., 16*
Gildersleeve, R. G., 147*, 1841
Gildersleeve, R. P., 60*, 110*, 167*
Gill, D. J., 1518
Gingher, P. E., 70*
Giraldo, C. A., 47*, 171*
Gjessing, K. M., 121*
Glahn, R. P., 90*, 1034, 1250, 1694
Gleaves, E. W., 91*
Glick, B., 16*, 33*, 114*, 135*, 855, 1210
Godke, R. A., 55*
Godwin, O., 45*
Goeger, M. P., 125*, 513
Golan, F. A., 1568
Gomez, F. A., 16*
Gomez-Gonzalez, F., 91*
Gonzales, L., 85*
Goodwin, M. A., 1694
Goodwin, T. L., 300
Gore, T. Z., 1632
Gorham, S. L., 730
Graves, H. B., 1008
Graves, J. G., 43*
Green, J., 17*, 1853
Greene, S. F., 95*
Grimes, J. L., 17*, 92*
Grossman, M., 33
Grumbles, L., 60*
Grunder, A. A., 183
Guenter, W., 98*, 147*
Guerra, C. M., 95*
Guise, K. S., 86*, 154*, 1794
Guodong, L., 92*
Gvaryahu, G., 399, 1642
Gyles, N. D., 40*
Gyles, N. R., 173*, 1123

H

Haddad, E. E., 92*, 93*
Hagan, F., 1210
Hagelberg, S., 707
Hagler, W. M., Jr., 101*
Hairston, H., 169*
Hakkinen, M., 1026
Halbrendt, C., 50*
Hall, R. D., 407
Hallett, J. S., 93*
Halvorson, D., 66*
Halvorson, J. C., 94*, 101*, 596
Hamilton Estiri, S., 75*
Hamilton, P. B., 58, 608, 619, 787, 1080
Hamm, D., 510
Hamre, M. L., 67*, 1306
Han, Y., 65*, 94*, 97*
Hancock, R. G., 91*
Hanners, G. D., 1166
Hansen, H. J., 98*
Hardin, J. M., 5*
Hargis, B. M., 79*, 95*
Hargis, P. S., 95*
Harms, R. H., 9*, 15*, 32*, 56*, 67*, 68*, 86*, 120*, 126*, 148*, 336, 452, 760, 794, 842, 1222, 1233
Harnish, S., 455
Harper, A. E., 1717
Harris, G. C., Jr., 2*, 17*, 18*, 23*, 48*, 135, 333, 1183
Harris, H. M., 10*
Harrison, P. C., 124*, 142*
Hart, I. C., 70*
Harter-Dennis, J. M., 87*, 96*, 1647
Harvey, R. B., 90*, 99*, 106*, 243, 253, 1139, 1418, 1778
Hashish, S., 84*
Hasiak, R. J., 585, 1062
Hattori, T., 1580
Havenstein, G. B., 108*, 149*, 166*, 191, 357, 1372, 1378, 1388, 1660, 1807
Heath, J. L., 33*, 122*, 165*, 1557, 1708
Hebrank, J. H., 110*
Heidelbaugh, N. D., 243
Heilman, L., 109*
Hellwig, H. M., 39*, 103, 170*
Hempe, J. M., 902
Henrikson, C. K., 101*
Henry, P. R., 18*, 41*, 1295, 1585
Hermes, J. C., 96*
Herro, D. R., 97*
Hess, J. B., 18*, 97*, 205
Hester, P. Y., 97*
Hickling, D., 98*
Hill, C. H., 174*
Hoerning, E. F., 123*
Hoes, T. L., 162*
Hoffbeck, S. L., 170*
Holbrook, D. M., 123*
Holleman, K. A., 98*
Holman, J. R., 53*
Holtzen, H., 41*
Hooper, M. J., 367
Houvenaghel, A., 1615
Howser, G. G., 871
Hoyle, C. M., 19*
Hsun, C. L., 115*
Huber, C. S., 138*, 139*
Hudson, H. A., 98*
Huerta-Leidenz, N., 91*
Huff, W. E., 90*, 99*, 106*, 253, 1034, 1139, 1418
Huffman, D. L., 54*
Huiatt, T. W., 87*
Hulan, H. W., 99*, 615, 1447, 1743
Hulet, R. M., 20*, 99*, 114*, 149
Hult, K., 707
Hunt, H., 126
Hunt, J. R., 156, 1020, 1243
Hurwitz, S., 384, 420, 1407
Hussain, N. A., 46*
Hussein, A. S., 19*, 41*, 62*, 137*, 149*, 1157
Huyghebaert, G., 950

I

Iciek, L., 88*
Imai, C., 261
Ingram, D. R., 1*, 19*, 35*
Isaacks, R. E., 100*
Ito, Y., 1290
Ivey, F. J., 12*, 37*, 76*, 1314
Izat, A. L., 10*, 13*, 38*, 43*, 69*, 80*, 133*, 1431, 1568
Izquierdo, O. A., 94*, 232, 341

J

Jabe, J. M., 100*
Jabe, M., 116*
Jackson, M., 111*
Jacques, K. A., 7*
Janky, D. M., 126*, 150*, 544, 715, 1222, 1545
Janni, K., 66*
Jayawickrama, L. H., 151*
Jaynes, J. M., 55*
Jeffers, T. K., 1043, 1050, 1058
Jenkins, M. C., 100*, 103*, 111*
Jensen, L. S., 28*, 119*, 139*, 280, 294, 1469
Jevne, C. J., 101*
Jiang, Z., 156*
Jin, L., 1400
John, T. M., 463
Johnson, A. D., 101*
Johnson, A. L., 101*, 165*
Johnson, J., 248
Johnson, M. G., 157*
Johnson, R. J., 118*
Johnson, T., 102*
Johnson, T. H., 7*, 19*, 41*, 62*, 137*, 149*, 1157
Johnson, W. A., 25, 32*, 55*
Johnson, Z., 1414
Johnson, Z. B., 39*, 103, 135, 333
Johnston, N. P., 173*
Johnstone, B. J., 152*
Jones, J. E., 24*, 115*
Jones, W. T., 102*, 137*
Julian, R. J., 1102

K

Kafri, I., 1356
Kaifer, C. R., 113*
Kakuda, Y., 270
Kamar, G. A., 102*, 113*
Kashani, A. B., 445
Kasivivat, K., 161*
Keene, O. D., 106*
Keirs, R., 17*, 1853
Keister, D. M., 72*

Kellogg, T., 17*
Kelly, B. J., 173*
Kennedy, T. J., 122*
Kerrigan,B., 1486
Keshavarz, K., 20*, 750
Khalili, A. D., 1593
Kido, S., 476
Kienholz, E. W., 138*
Kim, C. S., 72
Kim, J. W., 674
Kim, K. S., 103*
Kim, S., 1525
Kimmel, J. R., 126
Kimura, K., 1636
King, A. J., 103*, 134*
Kirby, J. D., 103*
Kirk, T. A., 20*
Klasing, K. C., 104*, 152*, 626
Klingensmith, P. M., 1203
Knight, C. D., 12*, 1314
Kocsis, J. F., 64*, 105*
Kogut, M. H., 105*
Kohl, H., 97*
Koike, T. I., 171*
Koldovsky, O., 153*
Kondo, F., 1424
Koops, W. J., 33
Kostelc, J. G., 76*
Kotula, K. L., 106*
Kouakou, B., 76*
Koyanagi, F., 1770
Kradel, D. C., 106*
Kratzer, F. H., 1818
Krause, G. F., 1455
Krautmann, B. A., 85*
Krieg, R., 455
Krista, L. M., 660
Krueger, W. F., 54*, 63*, 79*, 103, 133*
Kubena, L. F., 90*, 99*, 106*, 243, 253, 1139, 1418, 1778
Kuenzel, W. J., 107*
Kunkle, W. E., 29*

L

Lacy, M. P., 4*
Ladbrooke, B. D., 1731
Lalli, M. F., 819
Lamb, I. C., 107*
Lamont, S. J., 989
Lange, C., 105*
Larsen, C., 1229
Larsen, C. T., 99*, 149
Latimer, K. S., 1684
Latorre, J. R., 135, 333
Latshaw, J. D., 53*, 108*
Laurent, S. M., 1*, 15*, 18*, 19*, 35*
Laurin, D. E., 104*
Lauterio, T. J., 120
Lauxen, R. C., 902
Lawrence, L., 6*
Lawson, J. W., 26*
Laycock, S. R., 108*
Lee, H., 15*, 20*
Lee, H.-Y., 109*
Lee, M.-R., 34*, 156*
Lee, S. R., 21*, 109*
Lee, Y. B., 134*
Leenstra, F. R., 1, 10, 16
Leeson, S., 88, 391, 1102, 1680
Legendre, M. G., 123*
Leighton, Jr., A. T., 1505, 1518
Leske, K., 109*
Lev, Y., 809
Levenick, C. K., 1505
Lewis, R. H., 110*
Ley, D. H., 116*
Li, Q., 110*
Li, S., 111*
Lien, R. J., 21*, 110*
Lightsey, S. F., 24*, 30*, 115*, 144*
Lilburn, M. S., 21*, 22*, 51*, 161*
Lillard, H. S., 2*, 77*
Lillehoj, E. P., 111*, 1525
Lillehoj, H. S., 100*, 103*, 111*, 1525
Lin, C. Y., 565
Lin, H. K., 141*
Lin, S. Y., 112*
Lin, Y. C., 961
Link, G. A., 87*
Liu, J. K., 101*. 170*
Llames, C., 40*
Lomax, K. M., 113*
Long, D. W., 51*
Long, P. L., 248
Lott, B. D., 5*, 375, 677, 1239, 1514
Lowe, P. C., 914
Lucariello, M. A., 22*
Lukoszek, R. W., 110*
Luo, L., 21*
Lyon, B. G., 23*, 112*, 736
Lyon, C. E., 11*, 23*, 35*, 112*, 175*, 510, 736, 1549
Lyons, J. J., 49*, 94*, 164*, 407
Lyons, M. L., 851

M

Mabon, J., 168*
MacDonald, A., 85*
Macy, L. B., 17*, 18*, 23*
Mahmoud, A. Z., 102*, 113*
Maki, A. A., 113*
Mallinson, E. T., 730
Malone, G. W., 32*, 33*, 113*
Manley, R. W., 99*, 114*, 149
Mannig, D., 24*
March, B. E., 766
Marion, J. E., 29*, 544
Marks, H. L., 28*, 64*, 519, 1111, 1338
Marquardt, R. R., 147*, 502
Marquardt, W. W., 730
Marsh, J. A., 141*
Marshall, T. T., 29*
Marsh-Salin, J., 114*
Martin, F. G., 27*
Martin, P. W., 1684
Martosiswoyo, A. W., 280, 294
Mashaly, M. M., 83*, 92*, 93*, 114*, 166*, 1775
Maslin, W. R., 5*
Mast, M. G., 125*, 156*, 168*
Mastro, A. M., 1015
Masuda, S., 1770
Mathis, G. F., 115*
Maupin, T., 35*
Maurer, A. J., 44*, 102*, 1797
Maurice, D. V., 8*, 24*, 30*, 65*, 115*, 144*, 427
May, J. D., 25*, 58*, 64, 73*, 116*, 375, 1514
Maynard, P. M., 104*
McBride, M. A., 116*
McCombs, J. K., 1203
McCorkle, F. M., 75*, 88*, 100*, 116*
McCormack, J. F., 25*, 1874
McDaniel, G. R., 219, 660
McDonald-Jones, G., 117*
McDougald, L. R., 16*, 31*, 115*
McFarland, D. C., 121*
McFarlane, J. M., 1787, 1791
McGeachin, R. B., 52*, 130*
McGhee, G. C., 1001
McGinnis, J. P., 25*, 117*
McGuinness, M. C., 117*
McGuire, J., 219
McHan, F., 26*, 52*, 71*
McKeen, W. D., 118*
McKenzie, M. E., 248
McMillan, I., 57*, 1758, 1762
McMurtry, J. P., 61*, 97*, 118*, 146*, 1356
McNeill, J., 270
McRae. K. B., 1447
McRee, D. I., 60*
Meier, T. I., 26*
Meldrum, J. B., 149
Mellor, D. B., 75*
Melvin, V. K., 81*
Mench, J. A., 119*, 121*, 155*
Mendonca, C. X., Jr., 119*
Menken, B. Z., 1814
Merka, W. C., 119*
Merkley, J. W., 120*, 914
Metzler, M. J., 339
Meuten, D. J., 42*
Meyer, W. E., 120*
Mian, M., 15*
Miles, R. D., 15*, 18*, 41*, 68*, 86*, 120*, 345, 842, 1222, 1233, 1295, 1585
Millam, J. R., 120*
Miller, A. M., 79*
Miller, L., 121*, 378
Minshall, R. D., 121*
Misir, R., 1274
Mita, M., 261
Mitchell, G. E. Jr., 8*
Mobarak, M. S., 121*
Moeller, M. W., 122*
Mohamed-Ali, L. H., 122*
Mohammad, A. H., 122*
Mohammed, H. H., 1224*
Monsi, A., 327
Moody, N. W., 16*
Moore, R. W., 103

Moran, D. F., 39*
Moran, E. T., Jr., 123*, 144*, 463, 493, 1810
Moreng, R. E., 138*
Morgan, G. W., 19*, 378, 677
Mori, M., 1636
Mori, N., 1130
Morris, C. E., 123*
Morrison, W. D., 57*, 108*
Muir, W. M., 123*
Muiruri, H. K., 124*
Munn, B. J., 32*
Muramatsu, T., 1130
Murphy, B. D., 1062
Murphy, C. N., 1058
Murphy, D. W., 124*, 125*
Murray, D. L., 313, 669
Murray, W. W., 125*, 156*
Musa, R., 141
Myron, J. J., 36*

N

Naber, E. C., 74*, 160*, 455
Naito, H., 1290
Nakajime, T., 1580
Nakamaru, E., 261
Nakaue, H. S., 125*, 1619
Nam, C. H., 176*
Nauschuetz, W. F., 26*
Neiuwland, M.G.B., 79*
Nelson, C. E., 126*, 1225
Nelson, D. S., 126*
Nelson, J. R., 126*
Nestor, K. E., 22*, 144*, 149*, 166*, 357, 826, 1372, 1378, 1388, 1660, 1807
Nettleship, M. B., 1123
Newberry, R. C., 156, 1020, 1243
Newbrey, J. W., 590
Newcombe, M., 766
Newman, C. W., 1573
Newman, R. K., 1573
Nicastro, F., 7*
Nichols, C. R., 66*, 127*
Nichols, E. L., 10*
Nieberg, P. S., 173*, 367
Nielson, A., 162*
Niemann, L. M., 87*
Niemiec, J., 127*, 163*
Nieuwland, M. G. B., 79*
Nilipour, A. H., 1619
Nir, I., 140*
Nishiyama, 1770
Nissen, S., 87*
Nitsan, Z., 140*
Noguchi, T., 1290
Noll, S. L., 66*, 94*, 127*, 128*, 170*, 1750, 1760
Noormarji, S. H., 128*
Nordskog, A. W., 859
Norizzah, A. R., 128*
Nurmi, E., 1026
Nwokolo, E. N., 129*, 156*

O

O'Brien, B. A., 130*
O'Keefe, T. R., 1008
Obi, I., 1306
Obi, I. N., 129*
Obunike, J. C., 130*
Odom, T. W., 52*, 95*, 132*
Oduho, G. W., 1210
Ogasawara, D. M., 865
Oguntona, T., 141, 145
Oju, E. M., 130*, 170*, 1750, 1760
Okpokho, N. A., 848
Okumura, J., 1130
Olah, I., 16*
Omar, S., 130*, 131*
Ono, Y., 132*
Ontiveros, R. R., 76*
Opel, H., 141*
Orban, J. I., 27*, 132*
Ort, J. F., 17*, 67*, 92*, 101*
Ottinger, M. A., 110*, 121*, 1322
Ouart, M. D., 27*, 544, 1352
Owings, W. J., 132*, 585
Oyawoye, E. O., 133*
Oyofo, B. A., 133*

P

Palmer, D. H., 46*, 134*
Pancorbo, O. C., 3*
Pang, S. F., 66*
Pankratz, J., 152*
Papa, C. M., 11*, 23*, 25*, 117*, 275, 635, 1147
Papkoff, H., 145*
Pardue, S. L., 134*
Paredes, M. D. C., 1766
Parkhurst, C. R., 12*, 60*, 72*, 608
Parmer, E. L., 27*
Parsons, C. M., 65*, 94*, 97*, 341, 652
Patel, S. B., 134*
Patten, J. D., 18*, 1178
Patterson, D. L., 123*
Patterson, M., 95*
Patterson, P. H., 135*, 1329
Payne, C. J., 135*
Peebles, E. D., 28*
Peguri, A., 135*, 136*
Peng, R. K., 104*
Pescatore, A. J., 7*, 19*, 62*, 136*, 137*, 149*, 879, 1647
Pesti, G. M., 146*, 1188
Petersen, J., 1363
Peterson, R. A., 102*, 137*
Petitjean, M., 176*
Petitte, J. N., 117*, 137*
Petrowski, M. L., 95*
Pfaff, W. K., 138*
Philips, T. D., 106*, 243, 253
Phillips, R. E., 93*, 138*
Pike, O. A., 138*, 139*
Pimentel, J. L., 139*, 434
Pinchasov, Y., 28*, 139*, 140*
Pines, M., 809
Pit, R., 1, 10
Plavnik, I., 384, 420, 1407
Plumart, P. E., 140*
Pollock, H. G., 126
Porter, T. E., 95*
Poste, L. M., 680
Potchanakorn, M., 140*
Potter, L. M., 29*, 96, 99*, 114*, 140*, 1281
Presson, B., 40*, 173*
Price, J. F., 128*
Prichard, D. L., 29*, 1352
Primo, R. A., 103
Pro-M., A., 159*
Proudfoot, F. G., 615, 1447, 1743
Proudman, J. A., 141*
Pruett, S. B., 116*
Prusa, K. J., 1075
Puffenbarger, K. L., 35*
Puigserver, A., 56*, 81*
Purchase, H. G., 141*
Puzzi, J. V., 141*

Q

Quarles, C., 1*, 35*, 46*
Qureshi, M. A., 121*

R

Raharjo, Y. C., 153
Rahn, A. P., 76*, 142*
Rahn, B. E., 1089
Rangel-Lugo, M., 148*
Rao, K. S., 29*, 142*
Rao, P. V., 1295, 1585
Rauch, H.-W., 455
Raup, T. J., 3*, 30*
Raymond, L. N., 157*
Reagan, J. O., 11*, 23*
Reed, R. J., 9*, 128*
Reid, B., 153*
Reid, D. W., 1804
Reid, W. M., 1050
Reilly, W. M., 142*
Reine, A., 161*
Remsen, L. G., 143*
Remus, J. C., 143*
Renden, J. A., 34*, 143*, 219, 835, 897
Renner, P. A., 144*, 1660
Resurreccion, R. S., 1684
Revington, W. H., 144*, 1810
Reynells, R., 878
Reynolds, D. L., 132*
Rhee, K. C., 91*
Rhoades, J. F., 24*, 30*, 144*
Richardson, J. W., 89*
Riddell, C., 68*, 145*
Rikihisa, Y., 961
Riviere, J. E., 619
Robbins, K. R., 1001

Robel, E. J., 30*
Roberson, K. D., 1001
Roberson, R. H., 145*
Roberts, T., 936
Robey, W. W., 219, 800
Robinson, F. E., 68*, 145*, 153*, 1486
Robinzon, B., 157*, 1642
Rogers, B., 1236
Rogers, S. R., 146*
Roland, D. A., Jr., 1632
Roland, D. A., Sr., 27*, 29*, 31*, 132*, 142*, 577, 1801
Rolow, A., 1225
Rosebrough, R. W., 61*, 146*, 147*, 1356
Rosenberg, J., 809
Rotibi, A., 31*
Rotter, B. A., 147*
Rottinghaus, G. E., 1418
Rountree, J., 96
Rouse, J., 1225
Roush, W. B., 31*, 90*
Rowland, G. N., 6*, 150*
Rozenvoim, I., 1642
Ruff, M. D., 147*
Ruiz, N., 32*, 336, 148*, 760, 794, 1222
Ruiz-Lopez, B., 148*
Russek-Cohen, E., 730
Ruszler, P. L., 148*
Rutz, D. A., 871
Rutz, F., 149*
Rydell, J. A., 80*, 163*

S

Sacco, R. E., 144*, 149*, 1372, 1807
Saif, Y. M., 144*, 149*, 1372, 1660, 1807
Saito, F., 1130
Saito, J., 261
Salman, H. K., 85*
Salmon, R. E., 680, 1731
Sammelwitz, P. H., 149*
Sams, A. R., 88*, 150*
Sandel, G. W., 95*
Sanders, A. M., 150*
Sandusky, C. L., 1557, 1708
Sasamoto, S., 1625
Satterlee, D. G., 25, 32*, 60*, 313
Saunders, R. M., 1166
Savage, J. E., 159*, 902
Savage, P. K., 730
Savage, T. F., 55*, 1619
Sayag, N., 1642
Saylor, W. W., 32*, 33*, 113*, 151*
Scanes, C. G., 105*, 120, 470
Schaeffer, B. H., 52*
Schaeffer, J. L., 608, 619, 1080
Schally, A. V., 19*, 38*
Scheideler, S., 153*
Scheideler, S. E., 8*, 33*, 151*, 440
Scherer, G., 24*
Schieber, E. M., 1329
Schlegel, B. F., 1050, 1058
Schmieder, H., 43*
Schmisseur, E., 152*
Scoles, G. A., 871
Sconberg, S., 104*, 152*
Scott, H. M., 105*
Scott, T. R., 26*, 33*, 135*, 152*, 153*, 1210, 1781
Scott-Kline, D., 557
Sebranek, J. G., 1062
Seibert, B., 115*
Sell, J. L., 84*, 85*, 153*, 175*, 440, 585, 1089
Senkoylu, N., 154*
Sexton, K. J., 34*, 835, 897
Sexton, T. J., 131, 1483
Shackelford, A. D., 154*, 933, 1069
Shah, R. M., 865
Shapiro, R., 90*
Sharaf, M. M., 149*, 1372, 1807
Shaw, E. M., 154*
Shea, M. M., 119*, 155*
Shehata, M. A., 596
Sheldon, B. W., 57*, 73*, 155*
Shermer, W. D., 1725
Shippy, R. D., 50*
Shoffner, R. N., 154*
Shoop, M. A., 1783
Shultz, F. T., 173*
Shuman, R. M., 34*, 42*, 156*
Sibbald, I. R., 413, 1192
Siegel, H. S., 114*, 125*, 156*, 1008, 1775
Siegel, P. B., 800
Sieminski-Brodzina, L. M., 114*
Silsby, J. L., 86*, 93*, 95*, 128*, 138*, 1794
Sim, J. S., 129*, 156*
Simmons, D. G., 42*, 174*
Simmons, J. D., 34*, 64, 73*, 1239
Sims, M. D., 35*
Sinclair, R. W., 1486
Siopes, T. D., 21*, 110*, 1098
Siragusa, G. R., 157*
Skeeles, J. K., 40*, 135, 173*
Skeeles, K., 109*
Skewes, P. A., 114*
Skinner, J. T., 35*
Slajchert, T., 105*
Slater, T., 88*
Slaugh, B., 17*, 92*
Sloan, D. R., 27*
Slominski, B. A., 61*
Smalley, E. B., 174*
Smeltzer, M. A., 1684
Smith, C. 1655
Smith, D. B., 10*
Smith, D. M., 128*
Smith, D. P., 35*, 908, 1549, 1702
Smith, E., 191
Smith, E. J., 50*, 1668, 1674
Smith, M. O., 36*, 38*
Smith, N. K., 108
Smith, N. K., Jr., 108, 1174
Smith, O. E., 883
Smith, R. C., 1632
Smith, R. M., 56*
Smyth, J. R., Jr., 56*
Snapir, N., 157*, 1642
Sneed, L., 61*, 69*
Sneed, L. W., 158*
Snyder, D. B., 720, 730
Snyder, D. L., 1058
Soares, J. H., Jr., 82*, 1322
Soki, K., 1424
Somes, R. G., Jr., 158*
Sooncarernying, S., 159*
Sosa-M., E., 159*
Sosnicki, A., 973, 982
Southern, L. L., 47*, 51*, 171*, 172*
Spotts, C. B., 31*
Spurlock, M., 159*
Squires, M. W., 160*
Sreemannarayana, O., 502
St. Angelo, A. J., 123*
Stadelman, W. J., 36*, 43*, 160*
Stallings, K. K., 131*
Stamp, L. E., 51*
Stamp, L. M., 22*
Stamps, L., 2*, 48*, 1414
Stanger, N. E., 61*
Stanley, V. G., 161*
Steele, N. C., 61*, 146*, 147*, 1356
Steenson, L. R., 36*
Steffens, W. L., 1684
Steigner, J. W., 161*
Stephens, J. F., 55*, 74*, 160*, 689, 699
Stephenson, A. B., 1455
Stevens, V. I., 680, 1731
Stilborn, H. L., 36*, 37*, 161*, 1183
Stirtzinger, T., 1102
Strain, G. M., 143*
Strausberg, R. L., 72*
Strohlein, D. A., 100*, 147*
Strong, C. F., Jr., 54*, 162*
Suderman, D. R., 162*
Sugahara, M., 1580
Sulaiman, S., 24*, 115*
Sullivan, T. W., 80*, 163*
Summers, J. D., 88, 391
Sun, S. S., 1486
Sunde, M. L., 52*, 135*, 1329, 1771, 1829, 1717
Sundlof, S. F., 11*, 15*
Sutton, C. D., 219, 660
Sutton, R. W., 10*
Swann, G. S., 37*, 163*
Swatland, H. J., 1680
Swick, R. A., 12*, 37*, 76*, 1314
Swick, R. W., 135*
Swinkels, A. G. J. C., 79*
Szabo, J., 8*
Szymkiewicz, M., 127*, 163*

T

Tabib, Z., 58
Tadtiyanant, C., 164*
Tamas, T., 164*
Tamura, T., 814
Tasaki, I., 1130
Taya, K., 1625
Taylor, D. R., 243
Taylor, R. L., Jr., 75*, 88*, 100*,

116*, 164*, 1491
Teeter, R. G., 36*, 38*
Terao, H., 1290
Tesch, S. R., 165*
Thacker, H. L., 174*
Thaxton, J. P., 19*, 38*, 110*, 147*, 167*, 313, 669
Thayer, D. W., 1797
Thoma, J. A., 40*, 173*
Thomas, O. P., 85*, 1345
Thomas, R. A., 10*, 13*, 38*, 43*, 165*
Thompson, L. D., 715, 1545
Thompson, O. C., 74*
Tidwell, N. M., 39*
Tillman, P. B., 641, 647, 743
Tilly, J. L., 101*, 165*
Timmons, M. B., 534
Tivzenda, P. T., 6*
Toelle, V. D., 108*, 166*, 1378, 1388
Toensmeyer, U. C., 50*
Tonkinson, L. V., 1043, 1050, 1058
Tottori, J., 1424
Trammell, J. H., 103
Trout, J. M., 114*, 166*, 1775
Tsang, C. P. W., 166*, 1105
Tsiagbe, M. E., 1717
Tsiagbe, R. J., 1772
Tucker, R. E., 8*
Twining, P., 46*
Twining, P. V., Jr., 1*
Tyczkowski, J. K., 167*, 608, 619, 787, 1080

U

Ueberschaer, K.-H., 455
Uesugi, I., 261
Uijttenboogaart, T. G., 1536
Uskokovic, M., 166

V

Vagnoni, D. B., 172*
van der Zijpp, A. J., 79*
Van Krey, H. P., 20*, 114*, 800, 996
van Tienhoven, A., 399
Vandepopuliere, J. M., 49*, 58*, 59*, 94*, 164*, 407
Varghese, S. K., 167*
Vasilatos-Younken, R., 168*, 487, 826
Veit, H. P., 99*
Vena, V., 90*
Verrinder Gibbins, A. M., 70*, 92*
Vincent, S. C., 164*, 1491
Vo, K. V., 39*, 82*, 168*
Vogt, H., 455
Vohra, P., 1166
Voris, J. C., 169*

W

Wabeck, C. J., 33*
Wagner, D. D., 169*
Waibel, P. E., 94*, 101*, 127*, 130*, 170*, 596, 1750, 1760, 1814
Waldman, R., 160*
Waldroup, P. W., 4*, 6*, 7*, 13*, 18*, 22*, 36*, 37*, 38*, 39*, 103, 108, 161*, 165*, 170*, 300, 641, 647, 743, 1174, 1178, 1183, 1725
Wang, L., 69*
Wang, S., 30*, 171*
Ward, N. E., 40*
Ward, T. L., 171*
Ware, G. O., 64*, 1188
Washburn, K. W., 13*, 14*, 81*, 83*, 1781
Watkins, B. A., 45*, 171*, 307, 590, 778
Watkins, K. L., 47*, 51*, 171*, 172*
Weaver, W. D., Jr., 2*, 226, 996
Webb, J. E., 40*
Weber, H., 64*
Wechsung, E., 1615
Wedekind, K. J., 172*
Weinstock, D., 111*
Welch, C. C., 172*, 652
Wellenreiter, R. H., 168*
Wentworth, B. C., 93*, 1465
Whistler, P. E., 57*, 155*
White, G. W., 173*
White, W. B., 1329
Whitfill, C. E., 40*, 173*
Whiting, T., 48*
Wicker, D. L., 40*
Wideman, R. F., Jr., 90*, 107*, 1034, 1250, 1694
Widjajanti, P., 41*
Wierup, M., 1026
Wiggins, M., 1783
Wilke, B. N., 57*
Wilks, G., 164*
Williams, C., 37*
Williams, J. C., 143*
Wilmering, R. F., 685
Wilson, B. W., 173*, 367
Wilson, H. R., 15*, 56*, 336, 345, 685, 1233
Wilson, J. L., 660
Wilson, R., 510
Wineland, M. J., 63*
Winn, S. E., 53*
Winterfield, R. W., 174*
Witzel, D. A., 99*
Wold-Troell, M., 1026
Wolynetz, M. S., 413, 1192
Wong-Valle, J., 41*
Wood, N., 41*
Woodward, S. A., 150*, 715, 1222, 1264, 1545
Wu, W., 174*

X

Xia, L., 819

Y

Yagen, B., 787
Yamamoto, Y., 1210
Yates, J. D., 2*, 48*
Yazwinski, T. A., 41*
Yersin, A. G., 42*, 174*
Yoshimura, Y., 814
Young, D. C., 1058
Young, L. L., 175*, 510, 736
Youngren, O. M., 93*, 138*
Youssef-Hakimi, A. A., 513

Z

Zadworny, D., 319
Zarkadas, C. G., 1593
Zatari, I. M., 153*, 175*
Zeytun, I., 175*
Zhao, J., 42*, 156*
Zhiang, J., 129*
Zhu, D., 42*
Zimet, D. J., 1352
Zimmerman, N. G., 43, 47*, 126*, 176*
Ziprin, R. L., 176*
Zongo, D., 176*
Zubair, A. K., 141, 145
Zuckerman, A., 107*

Index of Subjects

1,25-dihydroxycholecalciferol, egg shell calcification, egg shell quality, layer performance, 1322

1,25-dihydroxyvitamin D_3, bone-breaking load, egg production, layer performance, plasma P, 1233†
egg shell quality, layer performance, vitamin D, 166*

17α-hydroxylase, hypophysectomy, ovarian follicle, quail-Japanese, theca cell, 1636

17β-estradiol dipropionate, lipase activity, pancreas, turkey poult, 84*

2,3-bisphosphoglycerate, erythrocytes, inositol pentakisphosphate, oxygen affinity, phosphate, 100*

3,3′,4,4′-tetrachlorobiphenyl, duck embryo, goose embryo, herring gull embryo, teratogenicity, 52

35 S-cystine, 35 S-methionine, egg albumen, sulfur amino acid level, 78*

35 S-methionine, 35 S-cystine, egg albumen, sulfur amino acid level, 78*

4-H, duck, fitting guidelines, showmanship guidelines, turkey, 167*

5-azacytidine, DNA methylation, gene expression, prolactin, turkey, 1794†
methylation, prolactin, turkey, 86*

6-hydroxydopamine, antibody production, broiler chick, catecholamines, 59*

Abdominal fat, adipocyte hyperplasia, adipocyte hypertrophy, caloric overconsumption, hyperplastic response, 800
adipose tissue, age, broiler, cellularity, 64*
age, broiler chick, feed efficiency, feed restriction, 384
bacitracin MD, body weight, broiler, iodine supplementation, 161*
beta agonist, eviscerated yield, feeding regimen, quail-Japanese, 120*
body weight, broiler sire strain, feed conversion, heritability, 1
selection, 10
breast yield, broiler, eviscerated carcass yield, feathering genotypes, 914
broiler, carcass measurement, feather meal, lysine, 111*
feed conversion, growth rate, light intensity, 73*
nutrient restriction, 6*
protein, 7*
selection, 65*
broiler chick performance, dressing percentage, energy feeding regimen, intestinal measurement, 131*
broiler growth, light intensity, 1239
broiler performance, dietary protein, 129*
feed restriction, glycolic acid, growth, 28*
meat and bone meal, metabolizable energy values, 294
metabolizable energy, poultry by-product meal, 119*
broiler sire strain, feed conversion, slaughter yield, weight gain, 16
genetic correlation, lipase, lipoprotein, protein, 183

Abdominal fat deposition, broiler, feather meal, finishing period, nonspecific nitrogen source, 300

Abdominal fat pad, adipocyte size, broiler, monensin, withdrawal period, 339†
am gene, broiler, eviscerated yield, 102*
broiler growth, broiler male chick, caponization, 979†
broiler performance, perlite, wheat, 33*
cloaca caliper measurment, turkey, 29*

Abnormal egg shell structure, calcium reserve, egg shell structure, 78*
palisade layer, 77*

Abnormal eggs, cage *vs.* floor housing, embryonic mortality, fertility, turkey, 1455

Accelerated rigor, electric stimulation, minimum time process, nonaged tenderness, tenderness, 40*

Accident rate, processing plant, 63*

Acetic and propionic acids, broiler chick, feed inhibition, feed intake, 139*

Acid water, broiler breeder, thyroid dysfunction, 109*

Acid-base balance, broiler, heat stress, nicarbazin, 3*
broiler chick, mineral anions, tibial dyschondroplasia, 148*
calcium level, pullet, urolithiasis, 90*
dietary calcium, dietary chloride, egg shell quality, sodium, 750
gypsum, layer diet, pulverized and particle size limestone, shell quality, 20*

Acid-base status, blood, heat stress, layer, thermoregulation, 113*
broiler growth, sodium, sodium bicarbonate, 151*

Acid-insoluble ash, amaranth, broiler, metabolizable energy, 641

Acquired immunity, broiler, *Cryptosporidium baileyi*, ELISA, immune response, 720

Acrosome reaction, cock spermatozoa, fertilization, *in vitro* incubation, perivitelline layer, 1770

Actin, avian skeletal muscle, collagen, myosin, N^{τ}-methylhistidine, 1593

Added fat in diet, quaifenesin in diet, reproductive performance, turkey hen, 114*

Adenohypophysis, age, chick, growth hormone, somatotrophs, 487

Adenosine triphosphate, electric stimulation, pH, poultry carcass, tenderness, 106*

Adipocyte hyperplasia, abdominal fat, adipocyte hypertrophy, caloric overconsumption, hyperplastic response, 800

Adipocyte hypertrophy, abdominal fat, adipocyte hyperplasia, caloric overconsumption, hyperplastic response, 800

Adipocyte size, abdominal fat pad, broiler, monensin, withdrawal period, 339†

Adipose cellularity, age, broiler growth, 1338

Adipose tissue, abdominal fat, age, broiler, cellularity, 64*
broiler pullet, energy source, feed access, 766

Adrenal, adrenocorticotropin, bursectomy, chick, *in vitro*, 83*

Adrenocortical cell subpopulation, corticosteroidogenesis, turkey, 105*

Adrenocortical cells, corticosteroidogenesis, cyclic adenosine monophosphate production, prostaglandin, 105*
corticosterone production, protein malnutrition, 64*

Adrenocorticotropin, adrenal, bursectomy, chick, *in vitro*, 83*
ascorbic acid, cell-mediated immunity, graft *vs.* host

*Indicates abstract.
†Indicates research note.

reaction, splenomegaly, 313
Adult selection, genetic merit, juvenile selection, layer performance, selection response, 1655
Aerobic fermentation, bacteria, broiler litter, 3*
broiler chick, oat hulls, poultry droppings, 52*
Aflatoxin, aluminas, aluminosilicate, detoxification, mycotoxin, 243
antibody response line, body weight, chick, chlortetracycline, 1229†
body weight, chick diet, hydrated sodium calcium aluminosilicate, 106*
broiler male chick, mycotoxin, synergy, T-2 toxin, 1418
carotenoids, lutein, pale bird syndrome, pharmacokinetics, 619, 1080
Age, abdominal fat, adipose tissue, broiler, cellularity, 64*
broiler chick, feed efficiency, feed restriction, 384
adenohypophysis, chick, growth hormone, somatotrophs, 487
adipose cellularity, broiler growth, 1338
beak trimming, feed consumption, guinea fowl, mortality, 141†
blood chemistry, broiler, mycotoxin, ochratoxin A, 1139
body weight, broiler male, season of the year, sudden death syndrome, 1243
breast muscle, broiler, protein degradation, protein synthesis, 135*
broiler, feed restriction, heat stress, mortality, 2*
diet, intestinal disaccharidases, turkey poult, 153*
egg yolk, layer, strain, 7*
growth hormone, metabolic clearance rate, protein restriction, secretion rate, 120
layer, methionine requirement, sulfur amino acid requirement, 170*
Age and weight-constant efficiency, broiler growth, feed efficiency, feed intake, weight correction, 565
Age at sexual maturity, dietary protein, dwarf broiler breeder performance, hatchability, reproductive performance, 219
Age constant, body weight, feed efficiency, heritability, weight constant, 1497
Age of bird, egg quality, egg storage position, shell treatment, yolk position, 75*
Aggressive behavior, broiler breeder male, feed restriction, tryptophan, 155*
Aging temperature, broiler breast meat, rigor development, 1147
Agonistic activity, cage density, dominance rank, egg production, layer, 198
Ahemeral lighting, egg production, egg weight, lighting program, pullet, 176*
Ahemeral photoperiod, egg production, laying photoperiod, rearing photoperiod, shell quality, 391
Air misting, broiler, evaporative cooling, 8*
Air sac lesions, broiler, feed efficiency, fructooligosaccharides, mortality, 46*
Air velocity, cage environment, energy consumption, feed intake, Leghorn cockerels, 62*
Alimet®, broiler chick, DL-methionine, excreta, metabolism, 12*
DL-methionine absorption, hindgut, 1314
brown egg layer, egg mass, feed efficiency, methionine, 16*
excreta, methionine, 76*
Allo-Kramer shear, anesthesia, biochemical values, broiler breast muscle, 25*
broiler breast meat, tenderness, texture profile analysis, 35*
texture, texture profile analysis, 1549
Aluminas, aflatoxin, aluminosilicate, detoxification, mycotoxin, 243
Aluminosilicate, aflatoxin, aluminas, detoxification, mycotoxin, 243
Aluminum, broiler, silicon, tibial dyschondroplasia, 84*
broiler chick, phosphorus metabolism, vitamin D, 19*
egg production, molt-induced, quail-Japanese, zinc, 1157
***Am* gene**, abdominal fat pad, broiler, eviscerated yield, 102*
Amaranth, acid-insoluble ash, broiler, metabolizable energy, 641
amino acid availability, broiler, 647
apparent metabolizable energy, broiler chick growth, nutritive value, 1166
body weight, broiler male performance, fat pad, processing yield, 743
Amino acid analysis, egg shell membranes, egg shell quality, shell-less egg, soft-shelled egg, 1203
Amino acid availability, amaranth, broiler, 647
Amino acid bioavailability, cecectomy, digestibility, poultry by-product meal, protein quality, 97*
Amino acid deficiency, arginine deficiency, chick growth, chloride excess, lysine deficiency, 172*
Amino acid(s), bioavailability, energy, guinea fowl, Leghorn, 145*
in vitro digestion, plant feedstuffs, true amino acid availability, 78
meat meal, protein quality, 65*
body weight, broiler male chick, feed conversion, threonine requirement, 108
broiler, carcass composition, lysine requirement, 161*
energy level, sex separate rearing, 39*
broiler male chick, tryptophan requirement, 1174
broiler performance, lysine, methionine, protein requirements, 36*
bushel weight, chick growth, corn, 21*
crude protein, regression equations, sorghum, 40*
D-amino acid oxidase, duckling, substrate specificity, 851†
fat, protein, turkey, 4*
growth rate, lysine, methionine, turkey tom performance, 86*
Ammonia, broiler production, litter treatment, 151*
Anamnestic response, antibody response, sheep red blood cells, 855†
Androgen, broiler, lighting program, organ weight, restricted feeding programs, 145*
Anesthesia, Allo-Kramer shear, biochemical values, broiler breast muscle, 25*
Aneuploids, bursal lymphocyte population, MHC genes, 74*
Animal by-product meal, metabolizable energy, protein, 72*
Animal fat, body weight, broiler chick performance, feed intake, monensin, 452
broiler performance, protein, 80*
Anorexia, energy metabolism, feed intake, thiamin deficiency, turkey, 143*
Anthelmintics, efficacy, nematodes, 41*
Antibiotic resistance, broiler carcass, market survey, *Salmonella*, 55*
Escherichia coli, food chain, kanamycin, *Salmonella*, 699
kanamycin, *Salmonella*, transconjugant, 689
Antibiotic treatment, glucose supplement, hatching metabolism, turkey embryo, 463
Antibiotics, antiurease antibody, chick growth, jackbean urease enzyme, passive immunity, 434

bacitracin, body weight, growth, guinea fowl, 145†
bovine luteinizing hormone, ovarian tissue, steroidogenesis, 1486†
broiler growth, small intestine, 43*
broiler performance, feed conversion, weight gain, 70*
semen diluent, sperm lysis, sperm motility, turkey, 131
Antibody, broiler chick, coccidia, *Eimeria tenella*, ELISA, 16*
Brucella abortus, heterophils, humoral immunity, male chickens, 1775†
immune response, layer, northern fowl mite, Western blot analysis, 549
Mycoplasma gallisepticum, *Mycoplasma synoviae*, 58*
Antibody level, coccidiosis, immunization, protection antigen, 147*
Antibody perturbations, chick, neural cell adhesion molecule, regenerating nerve fibers, 143*
Antibody production, 6-hydroxydopamine, broiler chick, catecholamines, 59*
environmental temperature, immune response, selection lines, sheep red blood cells, 79*
growth hormone, leukocyte distribution, lymphoid organs, T_3,T_4 doses, 92*
heat stress, immune response, selection lines, sheep red blood cells, 79*
immune response, light regimen, thymidine uptake, white blood cell, 114*
Antibody response, anamnestic response, sheep red blood cells, 855†
B-congenic chickens, *Eimeria acervulina*, p150 merozoite protein, recombinant coccidial antigens, 111*
egg production, Newcastle disease virus, *Pasteurella multocida*, turkey, 1372
histocompatibility complex, immunocompetence, phagocytosis, T cell-mediated response, 989
Pasteurella multocida, turkey, 149*
Antibody response line, aflatoxin, body weight, chick, chlortetracycline, 1229†
Anticoccidial drugs, broiler, coccidial lesion scores, growth performance, litter moisture, 126*
chick, monensin, protein level, protein source, 652
coccidiosis, halofuginone, nicarbazin, polyether ionophores, 31*
maduramicin, narasin, nicarbazin, salinomycin, 35*
Anticoccidial efficacy, battery cage trials, *Eimeria*, ionophorous antibiotic, narasin, 1043
BAY Vi 9142, coccidiosis, postinfection, toltrazuril, 122*
coccidiosis, diclazuril, *Eimeria*, 115*
Eimeria, floor pen trials, ionophorous antibiotic, narasin, 1050
field trials, monensin, narasin, 1058
Antifungal activity, gentian violet, heat, moisture, particle size, 58
Antimicrobial agents susceptibility, broiler, *Clostridium colinum*, ulcerative enteritis, 1424
Antinutritive effects, cecectomy, layer, rapeseed meal, 61*
Antioxidant, lipid oxidation, mechanically deboned meat, oxidative stability, 270
Antisperm antibodies, ELISA, immunohistochemistry, turkey fertility, 20*
Antitumor response, *B* complex genotype, histocompatibility complex, Rous sarcoma virus, 164*
Antiurease antibody, antibiotics, chick growth, jackbean urease enzyme, passive immunity, 434
Antiviral, infectious bronchitis virus, infectious bursal disease virus, 40*, 173*
Aorta, *Bordetella avium*, elastin, trachea, turkey poult, 174*
Apolipoprotein A-I, carotenoids, chick intestine, coccidiosis, *Eimeria acervulina*, 45*
Apparent metabolizable energy, amaranth, broiler chick growth, nutritive value, 1166
corn, soybean meal, true metabolizable energy, 10*
meat and bone meal, metabolizable energy, true metabolizable energy, 280
Arginine deficiency, amino acid deficiency, chick growth, chloride excess, lysine deficiency, 172*
Arginine utilization, chick, growth assay, lysine utilization, monensin, 341†
Arginine vasotocin, broiler, heat stress, mesotocin, surgery stress, 171*
Arprinocid, broiler resistance, *Eimeria tenella*, glycamide, nicarbazin, 164*
Artificial insemination, fertility, semen extender, spermatozoa-damaged, turkey semen, 1483†
Ascites, broiler, environmental temperature, roaster, 87*
Ascorbic acid, adrenocorticotropin, cell-mediated immunity, graft *vs.* host reaction, splenomegaly, 313
aspirin, broiler male chick growth, heat stress, 1183
broiler male chick, corticosterone implant, lipogenesis, 1356†
Ascorbic acid level, stress, turkey neonate, 134*
Ash, bomb calorimeter, feedstuffs, proximate analysis, 513†
Aspirin, ascorbic acid, broiler male chick growth, heat stress, 1183
Atherosclerosis, cholesterol, exercise, quail-Japanese, 161*
genetic selection, quail-Japanese, 51*
Attenuated vaccine, *Escherichia coli*, *Pasteurella multocida*, 113*
Attitudes of integrators, biosecurity, broiler industry survey, feed efficiency, growth regulators, 50*
Aureomycin, egg production, pelleting, pullet growth, sunflower seeds, 445
Australia, broiler, diseases, 89*
Avian leukosis virus, congenital transmission, endogenous virus, sex-linkage, slow-feathering, 1668
slow-feathering, tumors, 1674
early feathering, endogenous virus, genes, Z chromosome, 191
endogenous virus, hemagglutination, R_2 alloantisera, 50*
Avian skeletal muscle, actin, collagen, myosin, N^{τ}-methylhistidine, 1593
Avizyme®, barley, broiler diet, full-fat canola seed, layer diet, 156*
Rous sarcoma virus, tumor regression, 1491
***B*-actin gene**, gene mapping, *in situ* hybridization, metaphase chromosome, 154*
***B*-adrenergic agonist**, chick, euglycemic clamp technique, glucose, insulin, 470
cimaterol, duck, 73*
B-cells, growth hormone, natural killer cells, T-cells, T_3,T_4 doses, 93*
***B* complex genotype**, antitumor response, histocompatibility complex, Rous sarcoma virus, 164*
***B*-congenic chickens**, antibody response, *Eimeria acervulina*, p150 merozoite protein, recombinant coccidial antigens, 111*
flow cytometric analysis, Ia antigens, lymphocyte response, monoclonai antibody, 1525
B-endorphin, body temperature, feed intake, water intake, 25*
Bacitracin, antibiotic, body weight, growth, guinea fowl,

145†
body weight, broiler, narasin, roxarsone, 103
lasalocid, turkey, 99*
Bacitracin MD, abdominal fat, body weight, broiler, iodine supplementation, 161*
Bacteria, aerobic fermentation, broiler litter, 3*
broiler carcass speck count, corn, intestinal spillage, milo, 88*
Bacterial contamination, broiler carcass, cross-contamination, scalding-spay scalding, 77*
Bacterial flora, broiler meat, oxidative stability, storage time, 2*
Bactericide, lactic acid, poultry-processing water, *Salmonella*, 69*
Barley, Avizyme®, broiler diet, full-fat canola seed, layer diet, 156*
beta-glucanase, beta-glucans, broiler chick growth, Franubet cultivar, 1573
broiler, growth performance, 7*
chick performance, enzyme supplementation, viscosity, weight gain, 147*
corn texture, grit, pellets, turkey performance, 101*
egg weight, feed utilization, layer performance, metabolizable energy, 1306
Barley brewers grain meal, broiler performance, cholesterol, turkey performance, 175*
Barring, dermal melanin, linkage relationship, recessive white skin, sex chromosome, 530
Batter, broiler drumsticks, coating, egg albumen, 557
Battery cage trials, anticoccidial efficacy, *Eimeria*, ionophorous antibiotic, narasin, 1043
BAY Vi 9142, anticoccidial efficacy, coccidiosis, postinfection, toltrazuril, 122*
Beak trimming, age, feed consumption, guinea fowl, mortality, 141†
behavior, feed consumption, pullet, weight gain, 109*
body weight, broiler roaster growth, feed intake, pellets *vs.* mash, 1514
broiler chick performance, feed efficiency, heterophil:lymphocyte ratio, stress, 166*
Beef calves performance, broiler litter-corn diet, rye-ryegrass pasture, 29*
Beef sausage, fat content, meat flavor, quality attributes, turkey sausage, 1075
Behavior, beak trimming, feed consumption, pullet, weight gain, 109*
body weight, pullet chick, rearing density, 47*
broiler performance, light intensity, 1020
cage shape, feed trough partitions, layer performance, 1348†
caging system-experimental, chick performance, health, multiple stressors, 1787†
chick performance, electric shock-experimental system, health, stress, 1791†
Bermudagrass, egg production, egg weight, xanthophyll, yolk color, 1154
Beta agonist, abdominal fat, eviscerated yield, feeding regimen, quail-Japanese, 120*
Beta-glucanase, barley, beta-glucans, broiler chick growth, Franubet cultivar, 1573
broiler, growth performance, 7*
Beta-glucans, barley, beta-glucanase, broiler chick growth, Franubet cultivar, 1573
Bile acids, broiler chick, intestine, liver, SEHCAT, 17*
Biliary secretion, layer, pancreatic polypeptide, pancreatic secretion, 126
Bioassay, excreta collection, glucose, true metabolizable energy, water, 1192
Bioavailability, amino acid(s), energy, guinea fowl, Leghorn, 145*
in vitro digestion, plant feedstuffs, true amino acid availability, 78
meat meal, protein quality, 65*
body weight, phosphorus, toe ash, turkey, 96
broiler male chick, selenium, time, tissue uptake, 1295
chick, manganese, 41*
manganese methionine, 18*
selenium, tissue uptake, 1585
digestibility, methionine, methionine hydroxy analog, 94*
Biochemical values, Allo-Kramer shear, anesthesia, broiler breast muscle, 25*
Bioelectrical impedance, body composition prediction, broiler, 897
Bioelectrical impedance analysis, body composition, fat free mass, turkey breeder, 92*
Biogenic amines, duodenum, hypothalamus, mortality, turkey poult, 13*
Biological activity, immune response, interleukin-1, purification, 104*
Biosecurity, attitudes of integrators, broiler industry survey, feed efficiency, growth regulators, 50*
Biotin, broiler chick, fatty acids, phospholipids, 171*
chick, fatty acids, liver lipids, *trans*/18:1, 45*
Biotin bioavailability, canola meal, cereal grains, soybean meal, turkey poult, 1274
Biotin deficiency, bone modeling, broiler chick, linoleate, valgus-varus, 590
Biotin level, fertility, hatchability, turkey egg, 30*
Black light devices, caged layer, electrocuting flies, flies, muscalure, 871
Black-tailed red pattern, dark brown gene, gene interactions, plumage color, Villafranquina chickens, 351
Blackout rearing, body weight, broiler breeder performance, lighting program, 2*
Blended feed-grade fats, fatty acid composition, gas-liquid chromatography, *trans*/*cis* 18:1, 307
Blood, acid-base status, heat stress, layer, thermoregulation, 113*
Blood chemistry, age, broiler, mycotoxin, ochratoxin A, 1139
chick, metabolites, organ weight, sterigmatocystin, 502
formaldehyde, layer, pellet binder, urea, 37*
Blood leukocyte, dopamine, migration, norepinephrine, serotonin, 116*
Blood levels, calcium, layer, phosphorus, shell quality, 68*
Blood plasma, semen, seminal plasma, testosterone, turkey breeder, 1461
BOD removal, COD removal, hydrogen peroxide, poultry processing water, wastewater treatment, 24*
Body composition, bioelectrical impedance analysis, fat free mass, turkey breeder, 92*
broiler breeder, feed restriction, semen production, 34*
broiler breeder male, feeding regimen, semen characteristics, 835
compensatory growth, protein undernutrition, turkey performance, 1750
energy, protein, turkey breeder performance, 17*
Body composition prediction, bioelectrical impedance, broiler, 897
Body density, divergent selection, quail-Japanese, shank length, 88*
Body fat, chick, dietary energy, feed intake, meal feeding, 140*
Body size, body weight, fasting heat production, 146*
Body temperature, B-endorphin, feed intake, water intake,

25*
heat stress, naked neck *vs.* normal, 818
Body weight, abdominal fat, bacitracin MD, broiler, iodine supplementation, 161*
broiler sire strain, feed conversion, heritability, 1 selection, 10
aflatoxin, antibody response line, chick, chlortetracycline, 1229†
chick diet, hydrated sodium calcium aluminosilicate, 106*
age, broiler male, season of the year, sudden death syndrome, 1243
age constant, feed efficiency, heritability, weight constant, 1497
amaranth, broiler male performance, fat pad, processing yield, 743
amino acid(s), broiler male chick, feed conversion, threonine requirement, 108
animal fat, broiler chick performance, feed intake, monensin, 452
antibiotic, bacitracin, growth, guinea fowl, 145†
bacitracin, broiler, narasin, roxarsone, 103
lasalocid, turkey, 99*
beak trimming, broiler roaster growth, feed intake, pellets *vs.* mash, 1514
behavior, pullet chick, rearing density, 47*
bioavailability, phosphorus, toe ash, turkey, 96
blackout rearing, broiler breeder performance, lighting program, 2*
body size, fasting heat production, 146*
breast blister, broiler, cage floor material, 45*
broiler, feed conversion, wheat screenings, 615
probiotic, *Streptococcus faecium*, 132*
broiler breeder male, semen production, testes morphology, testis weight, 660
broiler breeder performance, egg production, lighting program, 48*
broiler breeder pullet, feed restriction, feeder space, 996
plasma corticosterone, plasma glucose, 809
broiler chick, feed conversion, soybean meal, tibial dyschondroplasia, 21*
broiler chick performance, ochratoxin A, organ weight, phenylalanine, 90*
broiler diet, litter condition, triticale, 1743
broiler male chick, deoxynivalenol, ochratoxin A, organ weight, 253
calcium, egg production, lead, quail-Japanese, 81*
carcass composition, genetic correlation, heritability, turkey, 1388
carcass yield, parts composition, turkey, 153*
chick growth, deoxynivalenol, organ weight, wheat contamination, 1778†
correlation, genetic selection, heat stress, 83*
dietary fat, feed efficiency, protein, turkey male, 1281
dwarf Leghorns, reproduction, sexual maturity, 143*
feed conversion, leg abnormalities, niacin requirement, turkey poult, 760
feed efficiency, feed intake, protein restriction, turkey male performance, 130*
feed restriction, molt, temperature, 137*
genetic correlation, growth hormone, heritability, plasma, 1781†
heritability, skeletal measurments, turkey, 1378
heat stress, naked neck, 13*
leg bones, leg muscles, shank width, turkey, 1660
lysine, threonine, turkey poult, valine, 140*
pancreas weight, soybean meal, turkey, urease index, 15*
salinomycin, turkey poult, viability, vomitoxin, 149†
turkey performance, virginiamycin, 94*
Body weight control, broiler breeder, monensin, phenypropanolamine, 133*
Body weight gain, broiler performance, Ethacal® feed component, temperature environment, 18*
Bomb calorimeter, ash, feedstuffs, proximate analysis, 513†
Bone, calcium level, egg shell weight, prelay pullet, 98*
egg production, egg shell quality, feed intake, magnesium excess, 97*
layer, magnesium excess, phosphorus, plasma, 18*
Bone ash, bone strength, bone structure, broiler performance, fluoride, 950
broiler, phosphorus availability, phosphorus sources, 152*
broiler chick, ionophores, magnesium, 59*
Bone calcium, chick, coccidiosis, feed efficiency, zeolite-A, 172*
Bone development, egg injection, embryo, vitamin A, vitamin D_3, 58*
Bone meal source, broiler performance, broiler strain, environmental temperature, phosphorus availability, 27*
Bone mineralization, broiler chick, hatching time, 59*
Bone modeling, biotin deficiency, broiler chick, linoleate, valgus-varus, 590
Bone strength, bone ash, bone structure, broiler performance, fluoride, 950
correlation, egg shell quality, 132*
Bone structure, bone ash, bone strength, broiler performance, fluoride, 950
Bone-breaking load, 1,25-dihydroxyvitamin D_3, egg production, layer performance, plasma P, 1233†
Boning time, broiler breast fillets, electric stimulation, postmortem metabolism, tenderness, 150*
Bordetella avium, aorta, elastin, trachea, turkey poult, 174*
cilia, collagen, tryptophan, turkey poult, 42*
Bovine luteinizing hormone, antibiotics, ovarian tissue, steroidogenesis, 1485†
Bovine serum albumin, chick, dopamine, ELISA, serotonin, 100*
Bovine serum albumin injection, host immunity, northern fowl mite, rocket electrophoresis, two-dimensional immunoelectrophoresis, 1015
Breast blister, body weight, broiler, cage floor material, 45*
broiler performance, feed efficiency, floor type, 48*
Breast muscle, age, broiler, protein degradation, protein synthesis, 135*
broiler, exercise patterns, muscle growth, thigh, 1708
cooking method, purification variables, thiamin content, 72
marination, moisture binding, rigor, shear force, 175*
muscle fiber diameter, muscle fiber type, 908
processing treatment, thigh muscle, turkey Canadian bacon, 162*
Breast up *vs.* breast down cooking, canola meal, cooking procedure, sensory quality, turkey, 680†
Breast yield, abdominal fat, broiler, eviscerated carcass yield, feathering genotypes, 914
Breed effect, nutrition, roaster weight, temperature regimen, 96*
Brewers dried grains, egg production, fertility, hatchability, pheasant breeder, 138*
Brine chilling, broiler breast fillets, cooking method, tenderness, 715
Brine solution *vs.* water, broiler breast fillets, chilling temperature, hot boning, tenderness, 1545

Broiler, abdominal fat, adipose tissue, age, cellularity, 64*
bacitracin MD, body weight, iodine supplementation, 161*
breast yield, eviscerated carcass yield, feathering genotypes, 914
carcass measurement, feather meal, lysine, 111*
feed conversion, growth rate, light intensity, 73*
nutrient restriction, 6*
protein, 7*
selection, 65*
abdominal fat deposition, feather meal, finishing period, nonspecific nitrogen source, 300
abdominal fat pad, adipocyte size, monensin, withdrawal period, 339†
am gene, eviscerated yield, 102*
acid-base balance, heat stress, nicarbazin, 3*
acid-insoluble ash, amaranth, metabolizable energy, 641
acquired immunity, *Cryptosporidium baileyi*, ELISA, immune response, 720
age, blood chemistry, mycotoxin, ochratoxin A, 1139
breast muscle, protein degradation, protein synthesis, 135*
feed restriction, heat stress, mortality, 2*
air misting, evaporative cooling, 8*
air sac lesions, feed efficiency, fructooligosaccharides, mortality, 46*
aluminum, silicon, tibial dyschondroplasia, 84*
amaranth, amino acid availability, 647
amino acid(s), carcass composition, lysine requirement, 161*
energy level, sex separate rearing, 39*
androgen, lighting program, organ weight, restricted feeding programs, 145*
anticoccidial drugs, coccidial lesion scores, growth performance, litter moisture, 126*
antimicrobial agents susceptibility, *Clostridium colinum*, ulcerative enteritis, 1424
arginine vasotocin, heat stress, mesotocin, surgery stress, 171*
ascites, environmental temperature, roaster, 87*
Australia, diseases, 89*
bacitracin, body weight, narasin, roxarsone, 103
barley, beta-glucanase, growth performance, 7*
bioelectrical impedance, body composition prediction, 897
body weight, breast blister, cage floor material, 45*
feed conversion, wheat screenings, 615
probiotic, *Streptococcus faecium*, 132*
bone ash, phosphorus availability, phosphorus sources, 152*
breast muscle, exercise patterns, muscle growth, thigh, 1708
carbonated water, feed intake, heat stress, mortality, 30*
carcass composition, carcass fat, fatty acids, 165*
carcass fat, plasma lipids, 129*
cellulose, crude fiber, water intake, 169*
chemical additives, microbial contamination, poultry-processing water, *Salmonella*, 10*
chiller water, diatomaceous earth, filtration, processing plant, 155*
chilling time, glycol solution, packaged carcasses, temperature, 154*
chlortetracycline, gentian violet, T-2 toxin, 99*
colonization, *Listeria monocytogenes*, 53*
compensatory growth, feed efficiency, feed restriction, 118*
feed restriction, growth hormone, 118*
composting, dead bird disposal, 124*
computer-controlled environment, heat resistance, heat stress, 12*
contract grower, economic analysis, rate of return, 1804†
corn, fatty acids, processing yield, sorghum, 130*
cryptosporidiosis, *Cryptosporidium* sp., cytologic preparations, 1684
digestive tract content, environmental temperature, feed regimen, 64
economics, layer, naked oats pricing, turkey, 10*
ELISA, monoclonal antibodies, *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, 116*
evaporative cooling, fan cooling, pad cooling, ventilation, 98*
feather release, microbiological contamination, picking feathers, spray scalding, 12*
feather release forces, scalding temperature, scalding time, stunning *vs.* nonstunning, 1069
feed efficiency, feed restriction, growth rate, sudden death syndrome, 1102†
pelleting, sunflower meal, 175*
feed intake, force feeding, plasma free fatty acids, plasma ketones, 30*
heat stress, water intake, 38*
plasma free arachidonic acid, 144*
feed restriction, heat stress, mortality, 226
growth hormone, insulin-like growth factor-1, plasma hormone level, 117*
heat stress resistance, 14*
hyperthermia, hypocalcemia, lactate, respiratory alklalosis, 132*
intestine, location, Meckel's diverticulum, yolk stalk, 677†
methionine, methionine hydroxy analog, oxidation, 81*
muscle tenderness, pH, post-mortem temperature, sarcomere length, 54*
prediction equation, water consumption, water:feed ratio, 131*
processing yield, vitamin C, 85*
Broiler breast fillets, boning time, electric stimulation, post-mortem metabolism, tenderness, 150*
brine chilling, cooking method, tenderness, 715
brine solution *vs.* water, chilling temperature, hot boning, tenderness, 1545
Broiler breast meat, aging temperature, rigor development, 1147
Allo-Kramer shear, tenderness, texture profile analysis, 35*
texture, texture profile analysis, 1549
broiler performance, lysine, methionine, 98*
denervation, lactate, pH, R value, 117*
electric stimulation, hot boning, poultry meat color, texture, 1536
electrical stunning, hot boning, meat quality, 674†
tenderness, textural quality, wing restraint, 275
Broiler breast muscle, Allo-Kramer shear, anesthesia, biochemical values, 25*
broiler drumsticks, broiler growth, exercise, muscle characteristics, 1557
cooking method, deboning time, post-mortem aging, texture profile, 23*
electric stimulation, texture profile, voltage levels, 23*
glycogen content, hot boning, lactate content, proximate analysis, 1702
R value, rigor development, sarcomere length, 635
Broiler breast tissue, fiber orientation, salt concentration, 122*
Broiler breeder, acid water, thyroid dysfunction, 109*
body composition, feed restriction, semen production,

34*
body weight control, monensin, phenypropanolamine, 133*
corticosterone, feed restriction, stress, tryptophan, 119*
dietary energy, quantitative traits, selection, 6*
divergent selection, fluorometry, sperm viability, 55*
egg production, feed restriction, sexual maturity, 1001
fat in diet, fertility, hatchability, lighting program, 5*
feed restriction, intestinal adaptation, 49*
feeding regimen, plasma calcium, plasma phosphorus, 56*
feeding time, hatchability, shell quality, slow *vs.* fast feathering, 538
Broiler breeder egg, fertility, hatchability, zeolite-A, 19*
Broiler breeder evaluation, glycerol, semen, tissue culture, 23*
Broiler breeder male, aggressive behavior, feed restriction, tryptophan, 155*
body composition, feeding regimen, semen characteristics, 835
body weight, semen production, testes morphology, testis weight, 660
diet, nitrogen retention, standard *vs.* all-corn diet, 144*
Broiler breeder performance, blackout rearing, body weight, lighting program, 2*
body weight, egg production, lighting program, 48*
egg production, fertility, hatchability, male-feeding system, 48*
floor type, hatchability, mortality, 1414
mortality, rabbit excreta, 39*
Ethacal® feed component, specific gravity, 35*
Broiler breeder pullet, body weight, feed restriction, feeder space, 996
plasma corticosterone, plasma glucose, 809
Broiler carcass, antibiotic resistance, market survey, *Salmonella*, 55*
bacterial contamination, cross-contamination, scalding-spay scalding, 77*
chemical preservative, feed, Myco Curb®, *Salmonella*, 1225†
detection, DNA probe, poultry processing water, *Salmonella*, 80*
ELISA, feed, gene probe, *Salmonella* isolation, 921
Broiler carcass quality, lysine, meat yield, 123*
Broiler carcass speck count, bacteria, corn, intestinal spillage, milo, 88*
Broiler chick, 6-hydroxydopamine, antibody production, catecholamines, 59*
abdominal fat, age, feed efficiency, feed restriction, 384
acetic and propionic acids, feed inhibition, feed intake, 139*
acid-base balance, mineral anions, tibial dyschondroplasia, 148*
aerobic fermentation, oat hulls, poultry droppings, 52*
Alimet®, DL-methionine, excreta, metabolism, 12*
DL-methionine absorption, hindgut, 1314
aluminum, phosphorus metabolism, vitamin D, 19*
antibody, coccidia, *Eimeria tenella*, ELISA, 16*
bile acids, intestine, liver, SEHCAT, 17*
biotin, fatty acids, phospholipids, 171*
biotin deficiency, bone modeling, linoleate, valgus-varus, 590
body weight, feed conversion, soybean meal, tibial dyschondroplasia, 21*
bone ash, ionophores, magnesium, 59*
bone mineralization, hatching time, 59*
carcass composition, compensatory growth, energy, feed restriction, 61*
colonization dose-50%, *Salmonella*, 71*
compensatory growth, energy repletion, enzyme activities, lipogenesis, 146*
corn-soybean meal diet, riboflavin requirement, 794
cortisol, cutaneous basophil hypersensitivity, mitogen-phytohemagglutinin, 669
DL-methionine, DL-methionine hydroxy analog, L-methionine, urinary excretion, 14*
delayed hypersensitivity, feed deprivation, feed intake-overconsumption, immunocompetence, 626
detection, enteropathogen, *Salmonella typhimurium*, 52*
digestibility, DL-methionine hydroxy analog, methionine, 14*
epiphyseal growth plate, growth hormone, tibial dyschondroplasia, 82*
Ethacal® feed component, sodium availability, 15*
flow rate, nipple waterer, 137*
glutathione peroxidase, glutathione reductase, riboflavin, selenium, 149*
roxarsone, selenium, 41*
hypersensitivity, immune response, soy protein, 104*
intestinal uptake, methionine hydroxy analog, rat, 56*
isoleucine, leg abnormalities, leucine, valine, 85*
lipogenesis, lysine, protein, 146*
magnesium deficiency, parathyroid, plasma minerals, 6*
protein solubility, soybean meal, urease activity, 11*
tibial dyschondroplasia, vitamin A, zeolite, 113
Broiler chick diet, by-product sources, phosphorus, precipitated bone phosphate, 1302
fat blends, poultry fat, yellow grease, 27*
Broiler chick growth, amaranth, apparent metabolizable energy, nutritive value, 1166
barley, beta-glucanase, beta-glucans, Franubet cultivar, 1573
feed intake, glycine, glycolic acid, glyoxylic acid, 1469
gizzard erosion, gizzerosine, mortality, 1580
Broiler chick hatch weight, dry bulb temperature, hatching time, wet bulb temperature, 163*
Broiler chick performance, abdominal fat, dressing percentage, energy feeding regimen, intestinal measurement, 131*
animal fat, body weight, feed intake, monensin, 452
beak trimming, feed efficiency, heterophil:lymphocyte ratio, stress, 166*
body weight, ochratoxin A, organ weight, phenylalanine, 90*
feed, supplemental equipment, water, 124*
layer performance, sesbania seed, 11*
management, semi-isolation brooding, weight gain, 96*
Broiler crosses, compensatory growth, energy restriction, feed efficiency, 156*
feather removal, feeding regimen, kosher processing, skin strength, 125*
Broiler diet, Avizyme®, barley, full-fat canola seed, layer diet, 156*
body weight, litter condition, triticale, 1743
calcium formate, formic acid, fumaric acid, organic acids, 1178
energy use, feed efficiency, light source, weight gain, 47*
feed additive, feed flavor enhancer efficacy, 22*
layer diet, Mazidagi raw rock phosphate, phosphorus source, 154*
soybeans, trypsin inhibitor, 37*
Broiler drumsticks, batter, coating, egg albumen, 557
broiler breast muscle, broiler growth, exercise, muscle characteristics, 1557

Broiler egg, immune response, incubation, microwave radiation, 57*
Broiler grower, interregional competition, profitability, simulation model, 89*
Broiler growout, broiler processing, competitive exclusion, *Salmonella* colonization control, 928
computer program, optimization, simulation program, temperature, 534
Broiler growth, abdominal fat, light intensity, 1239
abdominal fat pad, broiler male chick, caponization, 979†
acid-base status, sodium, sodium bicarbonate, 151*
adipose cellularity, age, 1338
age and weight-constant efficiency, feed efficiency, feed intake, weight correction, 565
antibiotics, small intestine, 43*
broiler breast muscle, broiler drumsticks, exercise, muscle characteristics, 1557
efficiency of production, light source, 151*
Broiler growth performance, florescent, incandescent, light source, sodium vapor, 43
Broiler industry survey, attitudes of integrators, biosecurity, feed efficiency, growth regulators, 50*
Broiler litter, aerobic fermentation, bacteria, 3*
computer simulation, economic value, litter feeding, 1352†
Broiler litter-corn diet, beef calves performance, rye-ryegrass pasture, 29*
Broiler male, age, body weight, season of the year, sudden death syndrome, 1243
estradiol, sudden death syndrome, 156†
feed efficiency, floor pen reared, fructooligosaccharides, 1*
Broiler male chick, abdominal fat pad, broiler growth, caponization, 979†
aflatoxin, mycotoxin, synergy, T-2 toxin, 1418
amino acid(s), body weight, feed conversion, threonine requirement, 108
tryptophan requirement, 1174
ascorbic acid, corticosterone implant, lipogenesis, 1356†
bioavailability, selenium, time, tissue uptake, 1295
body weight, deoxynivalenol, ochratoxin A, organ weight, 253
Broiler male chick growth, ascorbic acid, aspirin, heat stress, 1183
Broiler male performance, amaranth, body weight, fat pad, processing yield, 743
heat stress, Maxiban®, mortality, nicarbazin, 17*
Broiler meat, bacterial flora, oxidative stability, storage time, 2*
cooking yields, heat stress, moisture pick-up, 27*
microwave cooking, *Salmonella*, 71*
Broiler meat quality, carcass weepage, lipid stability, vitamin E, 38*
Broiler muscle, chilling, fiber type, rigor mortis, 150*
electric stimulation, ion exchange, lactate dehydrogenase, 11*
Broiler performance, abdominal fat, dietary protein, 129*
feed restriction, glycolic acid, growth, 28*
meat and bone meal, metabolizable energy values, 294
metabolizable energy, poultry by-product meal, 119*
abdominal fat pad, perlite, wheat, 33*
amino acid(s), lysine, methionine, protein requirements, 36*
animal fat, protein, 80*
antibiotic, feed conversion, weight gain, 70*
barley brewers grain meal, cholesterol, turkey performance, 175*
behavior, light intensity, 1020
body weight gain, Ethacal® feed component, temperature environment, 18*
bone ash, bone strength, bone structure, fluoride, 950
bone meal source, broiler strain, environmental temperature, phosphorus availability, 27*
breast blister, feed efficiency, floor type, 48*
broiler breast meat, lysine, methionine, 98*
broiler pigmentation, pigment source, xanthophyll, 544
carcass composition, isoproterenol, weight gain, 95*
chick weight, egg injection, zeolite-A, 1*
chlorine, water quality, waterers, 125*
coccidiosis, heat stress, narasin, nicarbazin, 248
coliforms, *Escherichia coli*, Luprosil, 13*
corn substitution, Iraq rations, local barley, 122*
cottonseed meal, soybean meal, 76*
cryptosporidiosis, *Cryptosporidium*, ELISA, 730
cyclic watering, feed intake, water intake, 378
ethoxyquin, feed preservative, peroxide, rancidity, 1725
fasting, heat stress, potassium chloride, weight gain, 36*
fat, grain sorghum, metabolizable energy, tannin content, 80*
feed restriction, lighting program, skeletal diseases, 68*
growout process, productivity, quality, 4*
halofuginone, maduramicin, monensin, nicarbazin, 32*
methionine, monensin, nicarbazin, salinomycin, 33*
methionine level, sodium sulfate, 159*
productivity, quality, 4*
Broiler pigmentation, broiler performance, pigment source, xanthophyll, 544
Broiler processing, broiler growout, competitive exclusion, *Salmonella* colonization control, 928
budgeting techniques, costs, small scale plants, 76*
Campylobacter jejuni contamination, 1568
chilling, evisceration, *Salmonella* control, scalding, 933
Broiler production, ammonia, litter treatment, 151*
commercial chicken industry, educational tool, table egg production, video, 134*
educational tool, table egg production, video, 46*
Broiler pullet, adipose tissue, energy source, feed access, 766
carcass composition, growth hormone, pulsatile, 168*
growth hormone, insulin, pulsatile, T_3,T_4, 168*
Broiler resistance, arprinocid, *Eimeria tenella*, glycamide, nicarbazin, 164*
Broiler roaster growth, beak trimming, body weight, feed intake, pellets *vs.* mash, 1514
Broiler sire strain, abdominal fat, body weight, feed conversion, heritability, 1
selection, 10
feed conversion, slaughter yield, weight gain, 16
Broiler starter diet, soybeans roasted *vs.* raw, soybeans-extruded, turkey starter diet, 163*
Broiler strain, bone meal source, broiler performance, environmental temperature, phosphorus availability, 27*
Broiler yield, dietary energy, virginiamycin, 1222
Broken-out eggs, polypropylene container, refrigerated storage, 138*
Broody, egg production, egg quality, progesterone, turkey, 72*
Broody hen management, prolactin, turkey, 128*
Brown egg layer, Alimet®, egg mass, feed efficiency, methionine, 16*
Brucella abortus, antibody, heterophils, humoral immunity, male chickens, 1775†
Budgeting techniques, broiler processing, costs, small

scale plants, 76*
Buffers, glycogenolysis, hormones, turkey liver explants, 147*
Bursa, cyclophosphamide, dendritic cell, spleen, 114*
dendritic cell, secretory cell, 16*
Bursal cell transfer, dopamine, immune response, plaque-forming cell transfer, sheep red blood cells, 88*
Bursal lymphocyte population, aneuploids, MHC genes, 74*
Bursectomy, adrenal, adrenocorticotropin, chick, *in vitro*, 83*
Bushel weight, amino acid(s), chick growth, corn, 21*
Business aspect of poultry industry, curriculum, poultry programs of the future, scientific aspect of poultry industry, student recruitment, 879
Busulfan, chemosterilant, quail-Japanese, 93*
Butylated hydroxyanisole, cysteine, lathyrism, quail-Japanese chick, toxicity, 153†
By-product sources, broiler chick diet, phosphorus, precipitated bone phosphate, 1302
Cage area, corticosterone, dominance rank, egg production, layer performance, 399
Cage density, agonistic activity, dominance rank, egg production, layer, 198
Cage environment, air velocity, energy consumption, feed intake, Leghorn cockerels, 62*
Cage floor material, body weight, breast blister, broiler, 45*
Cage shape, behavior, feed trough partitions, layer performance, 1348†
layer performance, nipple waterer, waterer location, waterers per cage, 89*
Cage type, feeder type, hen density, layer performance, 102*
Cage *vs.* floor housing, abnormal eggs, embryonic mortality, fertility, turkey, 1455
Caged layer, black light devices, electrocuting flies, flies, muscalure, 871
costs, culling procedures, livability, management tool, 148*
despotism, dominance, peck order, welfare, 1008
Caging system-experimental, behavior, chick performance, health, multiple stressors, 1787†
Calbindin, egg production, heat stress, layer, 52*
Calcitriol, layer, shell strength, 65*
Calcium, blood levels, layer, phosphorus, shell quality, 68*
body weight, egg production, lead, quail-Japanese, 81*
chloride, phosphorus, tibial dyschondroplasia, zeolite, 1436
layer, phosphorus absorption, phosphorus excretion, 440
Calcium excess, chick, manganese utilization, phosphorus excess, 172*
Calcium formate, broiler diet, formic acid, fumaric acid, organic acids, 1178
Calcium intake, feed intake, layer, self-select diets, 24*
Calcium level, acid-base balance, pullet, urolithiasis, 90*
bone, egg shell weight, prelay pullet, 98*
Calcium requirement, phosphorus requirement, rickets, turkey poult, 150*
Calcium reserve, abnormal egg shell structure, egg shell structure, 78*
palisade layer, 77*
egg shell, structural unit, 77*
Calcium source, egg production, limestone solubility, particle size, shell quality, 66*
layer digestive system, limestone, particle size, solubility, 142*
limestone, particle size, solubility, 29*
Caloric overconsumption, abdominal fat, adipocyte hyperplasia, adipocyte hypertrophy, hyperplastic response, 800
***Campylobacter jejuni* contamination**, broiler processing, 1568
egg-processing plant, egg products, 1431
Cannibalism, chronic coccidosis, congenital anomalies, ear, rabbits, 102*
Canola meal, biotin bioavailability, cereal grains, soybean meal, turkey poult, 1274
breast up *vs.* breast down cooking, cooking procedure, sensory quality, turkey, 680†
Canola seed, full-fat canola seed, pelleting, seed- processing methods, turkey, 1731
Canthaxanthin, chick, hydroxyechinenone, isozeaxanthin, metabolism, 787
Caponization, abdominal fat pad, broiler growth, broiler male chick, 979†
Carbon sources, coliforms, fermentation, manure-cage layer, *Salmonella*, 74*
Carbonated water, broiler, feed intake, heat stress, mortality, 30*
Carboxymethyl cellulose, fertility, semen extender, semen viscosity, 1218
Carcass composition, amino acid(s), broiler, lysine requirement, 161*
body weight, genetic correlation, heritability, turkey, 1388
broiler, carcass fat, fatty acids, 165*
broiler chick, compensatory growth, energy, feed restriction, 61*
broiler performance, isoproterenol, weight gain, 95*
broiler pullet, growth hormone, pulsatile, 168*
carcass yield, corn gluten feed, feed efficiency, turkey performance, 585
compensatory growth, feed restriction, turkey male, 1407
energy, layer, metabolizable energy utilization, temperature, 136*
energy estimation, lipid fraction, phospholipids, protein content, 413
Carcass fat, broiler, carcass composition, fatty acids, 165*
plasma lipids, 129*
dietary fat, feed efficiency, growth rate, turkey, 420
Carcass measurement, abdominal fat, broiler, feather meal, lysine, 111*
Carcass preservation, dead bird disposal, mortality, preservation acid, 113*
Carcass quality, compensatory growth, organ proportions, protein undernutrition, turkey, 1760
Carcass size, marinated broiler, moisture absorption, moisture retention, sodium pyrophosphate, 510†
Carcass weepage, broiler meat quality, lipid stability, vitamin E, 38*
Carcass yield, body weight, parts composition, turkey, 153*
carcass composition, corn gluten feed, feed efficiency, turkey performance, 585
guinea fowl, meat characteristics, 82*
Care of poultry, feed intake principles, management, 90*
Carotene, drying method, grass-clipping length, turf Bermudagrass, xanthophyll, 1644†
Carotenoid composition, egg, lutein, serum, yolk, 608
Carotenoids, aflatoxin, lutein, pale bird syndrome, pharmacokinetics, 619, 1080
apolipoprotein A-I, chick intestine, coccidiosis, *Eimeria acervulina*, 45*
Carrageenan, *in vitro*, mitogenesis, peripheral blood lym-

phocytes, phytohemagglutinin-M, 33*
Cassia obtusifolia seed, coffee weed, layer performance, nutrient fortification, 15*
Catecholamines, 6-hydroxydopamine, antibody production, broiler chick, 59*
chicken, feed intake, peptides, turkey, 74*
dopamine, epinephrine, norepinephrine, 75*
Ceca, mucin layer, *Salmonella* attachment, 26*
Cecectomy, amino acid bioavailability, digestibility, poultry by-product meal, protein quality, 97*
antinutritive effects, layer, rapeseed meal, 61*
Cell culture, glycerol, spermatozoa-chicken, 135
Cell wall antigen, enzyme immunoassay, *Listeria monocytogenes*, 157*
Cell-mediated immunity, adrenocorticotropin, ascorbic acid, graft *vs*. host reaction, splenomegaly, 313
Cellularity, abdominal fat, adipose tissue, age, broiler, 64*
Cellulose, broiler, crude fiber, water intake, 169*
Cereal grains, biotin bioavailability, canola meal, soybean meal, turkey poult, 1274
Chemical additives, broiler, microbial contamination, poultry-processing water, *Salmonella*, 10*
Chemical composition, culls, layer, mortality, 94*
layer diet, naked oats, weathering, 115*
Chemical denaturation, egg white, egg white protein, gelation, 53*
Chemical preservative, broiler carcass, feed, Myco Curb®, *Salmonella*, 1225†
Chemosterilant, busulfan, quail-Japanese, 93*
Chewiness, cooking method, deboning time, hardness, *pectoralis* muscle, 112*
Chick, adenohypophysis, age, growth hormone, somatotrophs, 487
adrenal, adrenocorticotropin, bursectomy, *in vitro*, 83*
aflatoxin, antibody response line, body weight, chlortetracycline, 1229†
antibody perturbations, neural cell adhesion molecule, regenerating nerve fibers, 143*
anticoccidial drugs, monensin, protein level, protein source, 652
arginine utilization, growth assay, lysine utilization, monensin, 341†
B-adrenergic agonist, euglycemic clamp technique, glucose, insulin, 470
bioavailability, manganese, 41*
manganese methionine, 18*
selenium, tissue uptake, 1585
biotin, fatty acids, liver lipids, *trans*/18:1, 45*
blood chemistry, metabolites, organ weight, sterigmatocystin, 502
body fat, dietary energy, feed intake, meal feeding, 140*
bone calcium, coccidiosis, feed efficiency, zeolite-A, 172*
bovine serum albumin, dopamine, ELISA, serotonin, 100*
calcium excess, manganese utilization, phosphorus excess, 172*
canthaxanthin, hydroxyechinenone, isozeaxanthin, metabolism, 787
copper, histidine, zinc, 47*
cyclic adenosine-3′,5′-monophosphate, gastric mucosal cells, gizzard erosion, gizzerosine, 1290
cyclosporine, immunosuppression, salmonellosis, 176*
feathers, fiber optics, sex-linked coloration, 1680
feed intake, hypothalamus, neuropeptide, 107*
immune response, zinc deficiency, 139*
infectious bronchitis virus, recombinant DNA, viral diagnosis, 69*
insulin, organ culture, pancreas, 101*
intestine, mannose sensitivity, receptors, *Salmonella typhimurium*, 133*
precocious puberty, semen production, tamoxifen, 1642†
Chick diet, aflatoxin, body weight, hydrated sodium calcium aluminosilicate, 106*
copper, peanut skins, weight gain, zinc, 51*
Chick embryo, chimera chick, embryonic cell transfer, 137*
cholesterol, growth hormone, insulin-like growth factor-1, very low density lipoprotein, 71*
leupeptin, proteolysis, skeletal muscle cell culture, 87*
mammalian embryo, shell-less culture, 55*
Chick growth, amino acid deficiency, arginine deficiency, chloride excess, lysine deficiency, 172*
amino acid(s), bushel weight, corn, 21*
antibiotics, antiurease antibody, jackbean urease enzyme, passive immunity, 434
body weight, deoxynivalenol, organ weight, wheat contamination, 1778†
Eimeria acervulina, feed efficiency, lysine deficiency, 232
Chick incubation, microwave radiation, reproductive performance, 57*
Chick intestine, apolipoprotein A-I, carotenoids, coccidiosis, *Eimeria acervulina*, 45*
Chick performance, barley, enzyme supplementation, viscosity, weight gain, 147*
behavior, caging system-experimental, health, multiple stressors, 1787†
electric shock-experimental system, health, stress, 1791†
Chick tissue, fatty acids, hydrogenated, *trans*/*cis* 18:1, 778
Chick weight, broiler performance, egg injection, zeolite-A, 1*
Chick weight gain, coccidiosis, duodenal pH, monensin, zeolite-A, 171*
Chicken embryo, gene transfer, retrovirus vector, tissue-specific expression, 34*
Chicken meat, high energy electrons, irradiation pasteurization, *Salmonella*, 165*
Chicken patties, frying conditions, lipid content, moisture content, shortening quality, 9*
off-flavor, reheating methods, sensory profiling, thiobarbituric acid, 112*
reheating methods, sensory evaluation, storage time, thiobarbituric acid values, 736
Chiller water, broiler, diatomaceous earth, filtration, processing plant, 155*
Chilling, broiler muscle, fiber type, rigor mortis, 150*
broiler processing, evisceration, *Salmonella* control, scalding, 933
Chilling temperature, brine solution *vs*. water, broiler breast fillets, hot boning, tenderness, 1545
Chilling time, broiler, glycol solution, packaged carcasses, temperature, 154*
Chimera chick, chick embryo, embryonic cell transfer, 137*
Chloramphenicol acetyltransferase, gene transfer, retrovirus vector, tissue-specific expression, 156*
Chlordecone, corticosterone, embryo, microwave radiation, quail-Japanese, 60*
Chlorella, egg production, layer performance, 84*
Chloride, calcium, phosphorus, tibial dyschondroplasia, zeolite, 1436
ion balance, layer diet, shell quality, sodium, 39*
Chloride excess, amino acid deficiency, arginine defi-

ciency, chick growth, lysine deficiency, 172*
Chloride level, layer, performance index, phosphorus level, potassium level, 131*
Chlorine, broiler performance, water quality, waterers, 125*
Chlortetracycline, aflatoxin, antibody response line, body weight, chick, 1229†
broiler, gentian violet, T-2 toxin, 99*
Cholesterol, atherosclerosis, exercise, quail-Japanese, 161*
genetic selection, quail-Japanese, 51*
barley brewers grain meal, broiler performance, turkey performance, 175*
chick embryo, growth hormone, insulin-like growth factor-1, very low density lipoprotein, 71*
egg, full-fat canola seed, omega-3 fatty acids, serum, 129*
herring meal, layer performance, omega-3 fatty acids, 99*
Choline, methionine, turkey poult, 9*
Choline supplementation, egg yolk, high performance liquid chromatography, methionine supplementation, phospholipids, 1717
Chronic coccidosis, cannibalism, congenital anomalies, ear, rabbits, 102*
Cilia, *Bordetella avium*, collagen, tryptophan, turkey poult, 42*
Cimaterol, *B*-adrenergic agonist, duck, 73*
Citrinin, diuresis, kidney function, manure moisture, ochratoxin A, 1034
ochratoxin A, pullet, renal function, 90*
Clearance kinetics, layer, narasin, 8*
Cloaca caliper measurment, abdominal fat pad, turkey, 29*
Cloning, *Escherichia coli*, *Mycoplasma gallisepticum*, 26*
growth hormone, nucleotide sequence, 107*
Clostridium, psittacine, ulcerative enteritis, 75*
Clostridium colinum, antimicrobial agents susceptibility, broiler, ulcerative enteritis, 1424
Coating, batter, broiler drumsticks, egg albumen, 557
Coccidia, antibody, broiler chick, *Eimeria tenella*, ELISA, 16*
Coccidial lesion scores, anticoccidial drugs, broiler, growth performance, litter moisture, 126*
Coccidiosis, antibody level, immunization, protection antigen, 147*
anticoccidial drugs, halofuginone, nicarbazin, polyether ionophores, 31*
anticoccidial efficacy, BAY Vi 9142, post infection, toltrazuril, 122*
diclazuril, *Eimeria*, 115*
apolipoprotein A-I, carotenoids, chick intestine, *Eimeria acervulina*, 45*
bone calcium, chick, feed efficiency, zeolite-A, 172*
broiler performance, heat stress, narasin, nicarbazin, 248
chick weight gain, duodenal pH, monensin, zeolite-A, 171*
Eimeria, immune response, recombinant coccidial antigens, 68*
Eimeria tenella, lymphokines, 105*
recombinant DNA, vaccine, 72*
ELISA, recombinant coccidial antigens, 103*
Cock spermatozoa, acrosome reaction, fertilization, *in vitro* incubation, perivitelline layer, 1770
COD removal, BOD removal, hydrogen peroxide, poultry-processing water, wastewater treatment, 24*
Coffee weed, *Cassia obtusifolia* seed, layer performance, nutrient fortification, 15*
Coliforms, broiler performance, *Escherichia coli*, Luprosil, 13*
carbon sources, fermentation, manure-cage layer, *Salmonella*, 74*
Collagen, actin, avian skeletal muscle, myosin, N$^{\tau}$-methylhistidine, 1593
Bordetella avium, cilia, tryptophan, turkey poult, 42*
Colonization, broiler, *Listeria monocytogenes*, 53*
Colonization dose-50%, broiler chick, *Salmonella*, 71*
Color-filtered light, growth, photoperiod, turkey performance, 1505
Commercial chicken industry, broiler production, educational tool, table egg production, video, 134*
Compensatory growth, body composition, protein undernutrition, turkey performance, 1750
broiler, feed efficiency, feed restriction, 118*
feed restriction, growth hormone, 118*
broiler chick, carcass composition, energy, feed restriction, 61*
energy repletion, enzyme activities, lipogenesis, 146*
broiler crosses, energy restriction, feed efficiency, 156*
carcass composition, feed restriction, turkey male, 1407
carcass quality, organ proportions, protein undernutrition, turkey, 1760
Competitive exclusion, broiler growout, broiler processing, *Salmonella* colonization control, 928
epidemiological evaluation, *Salmonella* control, 1026
Composition, functional properties, mechanically deboned meat, washing, 87*
kamaboko-like products, mechanically deboned meat, poultry meat color, yield, 112*
Composting, broiler, dead bird disposal, 124*
Computer controlled environment, broiler, heat resistance, heat stress, 12*
Computer data collection, egg shell thickness, electronic gauge performance, 1105†
Computer program, broiler growout, optimization, simulation program, temperature, 534
Computer simulation, broiler litter, economic value, litter feeding, 1352†
Computer software, expert decision support system, expert system technology, layer management, 152*
Computer spreadsheet templates, contract, financial attractiveness, turkey growout, 142*
Computerized contest scoring, judging contest, scan sheet scoring, 140*
Computerized data collection system, egg shell quality, egg specific gravity, egg weight, 902
Conductance, egg shell permeability, hatchability, turkey, 9*
Conductive heat loss, foot temperature, heat stress, layer, thermal perch, 142*
heat stress, metabolic rate, panting, roost temperature, 124*
Confounding experimental results, energy supplementation, growth assay, oil, sand, 169*
Congenital anomalies, cannibalism, chronic coccidosis, ear, rabbits, 102*
Congenital transmission, avian leukosis virus, endogenous virus, sex-linkage, slow-feathering, 1668
slow-feathering, tumors, 1674
Congo Red negative, Congo Red positive, *Escherichia coli*, macrophages, 121*
Congo Red positive, Congo Red negative, *Escherichia coli*, macrophages, 121*
Constitution and by-laws, Poultry Science Association,

161
Consumer preference, consumer preference, package size, poultry products, 50*, 50*
fatty material injection, roasted turkey, turkey broth injection, 160*
Contract, computer spreadsheet templates, financial attractiveness, turkey growout, 142*
Contract grower, broiler, economic analysis, rate of return, 1804†
Control populations, egg production, egg weight, hatchability, quail-Japanese, 357
Controlled feeding, egg production, energy utilization, 136*
Cooked chicken, ELISA, *Listeria monocytogenes*, 157*
Cooked gels, mechanically deboned meat, spray-dried egg white, water-washed, 73*
Cooking, meat composition, meat source, turkey roast, 1810†
Cooking method, breast muscle, purification variables, thiamin content, 72
brine chilling, broiler breast fillets, tenderness, 715
broiler breast muscle, deboning time, post-mortem aging, texture profile, 23*
chewiness, deboning time, hardness, *pectoralis* muscle, 112*
marinated chicken fillet, moisture content, volatile analysis, yield, 128*
Cooking procedure, breast up *vs.* breast down cooking, canola meal, sensory quality, turkey, 680†
Cooking time, oven type, turkey roast size, 139*
Cooking yields, broiler meat, heat stress, moisture pick-up, 27*
Copper, chick, histidine, zinc, 47*
chick diet, peanut skins, weight gain, zinc, 51*
dyschondroplasia, fusarochromanone, immunity, mycotoxin, 174*
Corn, amino acid(s), bushel weight, chick growth, 21*
apparent metabolizable energy, soybean meal, true metabolizable energy, 10*
bacteria, broiler carcass speck count, intestinal spillage, milo, 88*
broiler, fatty acids, processing yield, sorghum, 130*
Corn gluten feed, carcass composition, carcass yield, feed efficiency, turkey performance, 585
Corn oil, egg mass, linoleic acid, 118*
Corn substitution, broiler performance, Iraq rations, local barley, 122*
Corn texture, barley, grit, pellets, turkey performance, 101*
Corn-soy-gelatin basal diet, methionine activity, methionine assay, soybean meal, 88
Corn-soybean meal diet, broiler chick, riboflavin requirement, 794
fertility, hatchability, niacin supplementation, turkey breeder performance, 336†
Correlation, body weight, genetic selection, heat stress, 83*
bone strength, egg shell quality, 132*
Corticosteroidogenesis, adrenocortical cell subpopulation, turkey, 105*
adrenocortical cells, cyclic adenosine monophosphate production, prostaglandin, 105*
Corticosterone, broiler breeder, feed restriction, stress, tryptophan, 119*
cage area, dominance rank, egg production, layer performance, 399
chlordecone, embryo, microwave radiation, quail-Japanese, 60*
feed restriction, quail-Japanese, testosterone, 121*
heritability, quail-Japanese, stress, 25
prostaglandin-E_2, quail-Japanese, stress, 32*
Corticosterone implant, ascorbic acid, broiler male chick, lipogenesis, 1356†
Corticosterone production, adrenocortical cells, protein malnutrition, 64*
Cortisol, broiler chick, cutaneous basophil hypersensitivity, mitogenphytohemagglutinin, 669
Costs, broiler processing, budgeting techniques, small scale plants, 76*
caged layer, culling procedures, livability, management tool, 148*
Cottonseed meal, broiler performance, soybean meal, 76*
Coturnix (see quail)
Coturnix International, educational program, goals and strategy, youth development, 167*
Cross-contamination, bacterial contamination, broiler carcass, scalding-spay scalding, 77*
Crude fiber, broiler, cellulose, water intake, 169*
Crude protein, amino acid(s), regression equations, sorghum, 40*
Cryptosporidiosis, broiler, Cryptosporidium baileyi, cytologic preparations, 1684
broiler performance, *Cryptosporidium*, ELISA, 730
Cryptosporidium, broiler performance, cryptosporidiosis, ELISA, 730
Cryptosporidium baileyi, acquired immunity, broiler, ELISA, immune response, 720
Cryptosporidium baileyi, broiler, cryptosporidiosis, cytologic preparations, 1684
Culling procedures, caged layer, costs, livability, management tool, 148*
Culls, chemical composition, layer, mortality, 94*
Curriculum, business aspect of poultry industry, poultry programs of the future, scientific aspect of poultry industry, student recruitment, 879
Cutaneous basophil hypersensitivity, broiler chick, cortisol, mitogen phytohemagglutinin, 669
Cyclic adenosine monophosphate, adrenocortical cells, corticosteroidogenesis, prostaglandin, 105*
chick, gastric mucosal cells, gizzard erosion, gizzerosine, 1290
diacylglycerol, luteinizing hormone-releasing hormone, luteinizing hormone secretion, 101*
Cyclic watering, broiler performance, feed intake, water intake, 378
Cyclophosphamide, bursa, dendritic cell, spleen, 114*
Cyclosporine, chick, immunosuppression, salmonellosis, 176*
Cysteine, butylated hydroxyanisole, lathyrism, quail-Japanese chick, toxicity, 153†
Cystine, layer performance, methionine, transsulfuration enzyme activity, 78*
Cytochrome c, heme-complex, ligand, myoglobin, poultry meat color, 44*
Cytogenetics, egg storage, embryo abnormalities, heteroploid, karyotype, 96*
Cytologic preparations, broiler, cryptosporidiosis, *Cryptosporidium baileyi* sp., 1684
Cytotoxic T cells, lymphocyte antigens, monoclonal antibodies, natural killer cells, 111*
D-amino acid oxidase, amino acid(s), duckling, substrate specifity, 851†
Dark brown gene, black-tailed red pattern, gene interactions, plumage color, Villafranquina chickens, 351
Dark turkey meat, protein, sodium salts, storage, 103*
sodium salts, storage time, water added, 134*

Day of incubation, egg turning, hatchability, high density plastic egg flat, malposition, 685
Days without eggs, fasting periods, feather loss, layer performance, peaks of production, 67*
Dead bird disposal, broiler, composting, 124*
carcass preservation, mortality, preservation acid, 113*
Deboning time, broiler breast muscle, cooking method, post-mortem aging, texture profile, 23*
chewiness, cooking method, hardness, *pectoralis* muscle, 112*
Defective eggs, genetic associations, oviposition rate, shell quality, shell-less egg, 123*
Dehulling, meat flavor, turkey growth, white lupine, 596
Delayed glycolysis, electric stunning, muscle pH, protein solubility, turkey muscle, 1062
Delayed hypersensitivity, broiler chick, feed deprivation, feed intake-overconsumption, immunocompetence, 626
Dendritic cell, bursa, cyclophosphamide, spleen, 114*
secretory cell, 16*
Denervation, broiler breast meat, lactate, pH, R value, 117*
pectoralis muscle, 60*
Deoxynivalenol, body weight, broiler male chick, ochratoxin A, organ weight, 253
chick growth, organ weight, wheat contamination, 1778†
Dermal melanin, barring, linkage relationship, recessive white skin, sex chromosome, 530
Despotism, caged layer, dominance, peck order, welfare, 1008
Detection, broiler carcass, DNA probe, poultry processing water, *Salmonella*, 80*
broiler chick, enteropathogen, *Salmonella typhimurium*, 52*
Detoxification, aflatoxin, aluminas, aluminosilicate, mycotoxin, 243
Diacylglycerol, cyclic adenosine monophosphate, luteinizing hormone releasing hormone, luteinizing hormone secretion, 101*
Diaminopimelic acid, lysine, microorganisms, rooster-intact, cecectomized or cannulated, 1345†
Diatomaceous earth, broiler, chiller water, filtration, processing plant, 155*
Diclazuril, anticoccidial efficacy, coccidiosis, *Eimeria*, 115*
Diet, age, intestinal disaccharidases, turkey poult, 153*
broiler breeder male, nitrogen retention, standard *vs.* all-corn diet, 144*
Dietary acidification, dietary alkalinization, layer performance, renal function, urolithiasis, 1694
Dietary alkalinization, dietary acidification, layer performance, renal function, urolithiasis, 1694
Dietary calcium, acid-base balance, dietary chloride, egg shell quality, sodium, 750
infectious bronchitis virus, kidney function, pullet chick, urolithiasis, 1250
Dietary chloride, acid-base balance, dietary calcium, egg shell quality, sodium, 750
Dietary energy, body fat, chick, feed intake, meal feeding, 140*
broiler breeder, quantitative traits, selection, 6*
broiler yield, virginiamycin, 1222
Dietary fat, body weight, feed efficiency, protein, turkey male, 1281
carcass fat, feed efficiency, growth rate, turkey, 420
hepatic tocopherol concentration, liver, turkey, vitamin E, 1814†
Dietary level, egg production, Haugh units, wheat middlings, 1329
Dietary protein, abdominal fat, broiler performance, 129*
age at sexual maturity, dwarf broiler breeder performance, hatchability, reproductive performance, 219
egg production, layer performance, pullet dietary treatment, 62*
energy, reproductive performance, triticale, turkey semen, 1619
plasma thyroxine, quail-Japanese, thiouracil, 28*
Dietary protein restriction, fowl cholera, humoral immunity, *Pasteurella multocida*, 135*
Dietary pyruvic acid, egg production, fatty liver hemorrhagic syndrome, feed intake, heat stress, 104*
Digestibility, amino acid bioavailability, cecectomy, poultry by-product meal, protein quality, 97*
bioavailability, methionine, methionine hydroxy analog, 94*
broiler chick, DL-methionine hydroxy analog, methionine, 14*
oligosaccharides, soybean meal, transit time, 109*
Digestive tract content, broiler, environmental temperature, feed regimen, 64
Dimethyl sulfide, egg products, odor, spoilage, uracil, 123*
Disease challenge, egg production, *Pasteurella multocida*, turkey, 1807†
Diseases, Australia, broiler, 89*
Diuresis, citrinin, kidney function, manure moisture, ochratoxin A, 1034
Diurnal rhythm, feather condition, layer, lighting program, operant heat demand, 108*
Divergent selection, body density, quail-Japanese, shank length, 88*
broiler breeder, fluorometry, sperm viability, 55*
DL-methionine, Alimet®, broiler chick, excreta, metabolism, 12*
broiler chick, DL-methionine hydroxy analog, L-methionine, urinary excretion, 14*
DL-methionine absorption, Alimet®, broiler chick, hindgut, 1314
DL-methionine hydroxy analog, broiler chick, D-methionine, L-methionine, urinary excretion, 14*
digestibility, methionine, 14*
DNA fingerprinting, relatedness between individuals, 92*
DNA methylation, 5-azacytidine, gene expression, prolactin, turkey, 1794†
DNA modifications, egg yolk protein precursor, quail-Japanese, vitellogenin gene, 70*
DNA probe, broiler carcass, detection, poultry processing water, *Salmonella*, 80*
Dominance, caged layer, despotism, peck order, welfare, 1008
Dominance rank, agonistic activity, cage density, egg production, layer, 198
cage area, corticosterone, egg production, layer performance, 399
Dopamine, blood leukocyte, migration, norepinephrine, serotonin, 116*
bovine serum albumin, chick, ELISA, serotonin, 100*
bursal cell transfer, immune response, plaque-forming cell transfer, sheep red blood cells, 88*
catecholamines, epinephrine, norepinephrine, 75*
Double oviduct, experimental surgery, fertility, oviposition, 1465
Dressing percentage, abdominal fat, broiler chick performance, energy feeding regimen, intestinal measurement, 131*
D-Trp-6-LHRH, gonadotroph, layer, molt, reproductive

performance, 38*
layer performance, molt-induced, 19*
Dry bulb temperature, broiler chick hatch weight, hatching time, wet bulb temperature, 163*
hatch weight, wet bulb temperature, 37*
Dry feed, egg production, heat stress, wet feed, 164*
Drying method, carotene, grass-clipping length, turf Bermudagrass, xanthophyll, 1644†
Duck, 4-H, fitting guidelines, showmanship guidelines, turkey, 167*
B-adrenergic agonist, cimaterol, 73*
Duck embryo, 3,3′,4,4′-tetrachlorobiphenyl, goose embryo, herring gull embryo, teratogenicity, 52
Duckling, amino acid(s), D-amino acid oxidase, substrate specificity, 851†
Duodenal pH, chick weight gain, coccidiosis, monensin, zeolite-A, 171*
Duodenum, biogenic amines, hypothalamus, mortality, turkey poult, 13*
feed consumption, layer, molt-induced, zinc, 58*
Dust-bathing activity, egg production, egg quality, fear-related response, quail-Japanese, 1363
Dwarf broiler breeder performance, age at sexual maturity, dietary protein, hatchability, reproductive performance, 219
Dwarf Leghorns, body weight, reproduction, sexual maturity, 143*
Dyschondroplasia, copper, fusarochromanone, immunity, mycotoxin, 174*
Ear, cannibalism, chronic coccidiosis, congenital anomalies, rabbits, 102*
Early feathering, avian leukosis virus, endogenous virus, genes, Z chromosome, 191
Early protein restriction, egg type pullet, growth restriction, layer performance, 54*
Economic analysis, broiler, contract grower, rate of return, 1804†
Economic costs, foodborne disease costs, risk assessment, *Salmonella* control, salmonellosis, 936
Economic development, extension poultry specialist role, 136*
Economic impact, egg breakage, shell quality, uncollectable eggs, 1801†
Economic value, broiler litter, computer simulation, litter feeding, 1352†
Economics, broiler, layer, naked oats pricing, turkey, 10*
Educational poultry display, teaching tool quiz, 63*
Educational program, Coturnix International, goals and strategy, youth development, 167*
Educational slides, egg grading and packing, egg- processing plant design, egg-processing slides, extension program, 54*
Educational tool, broiler production, commercial chicken industry, table egg production, video, 134*
table egg production, video, 46*
Efficacy, anthelmintics, nematodes, 41*
Efficiency of production, broiler growth, light source, 151*
Egg, carotenoid composition, lutein, serum, yolk, 608
cholesterol, full-fat canola seed, omega-3 fatty acids, serum, 129*
glycoprotein, membrane layers, ovomucin, vitelline membrane, 476
internal quality, storage condition, 133*
ochratoxin A, quail-Japanese, tissue distribution, whole body autoradiography, 707
Egg albumen, 35 S-cystine, 35 S-methionine, sulfur amino acid level, 78*
batter, broiler drumsticks, coating, 557
Egg assay for riboflavin, layer, nutritional status, riboflavin, 160*
Egg breakage, economic impact, shell quality, uncollectable eggs, 1801†
egg processing plant, shell quality measurements, 162*
Egg grading and packing, educational slides, egg- processing plant design, egg-processing slides, extension program, 54*
Egg injection, bone development, embryo, vitamin A, vitamin D_3, 58*
broiler performance, chick weight, zeolite-A, 1*
Egg mass, Alimet®, brown egg layer, feed efficiency, methionine, 16*
corn oil, linoleic acid, 118*
egg production, genetic parameters, selection index, shell strength, 42*
layer, lysine assay, soybean meal, 49*
Egg-processing plant, *Campylobacter jejuni* contamination, egg products, 1431
egg breakage, shell quality measurements, 162*
Egg-processing plant design, educational slides, egg grading and packing, egg-processing slides, extension program, 54*
Egg-processing slides, educational slides, egg grading and packing, egg-processing plant design, extension program, 54*
Egg production, (see layer performance also)
1,25-dihydroxyvitamin D_3, bone-breaking load, layer performance, plasma P, 1233†
agonistic activity, cage density, dominance rank, layer, 198
ahemeral lighting, egg weight, lighting program, pullet, 176*
ahemeral photoperiod, laying photoperiod, rearing photoperiod, shell quality, 391
aluminum, molt-induced, quail-Japanese, zinc, 1157
antibody response, Newcastle disease virus, *Pasteurella multocida*, turkey, 1372
Aureomycin, pelleting, pullet growth, sunflower seeds, 445
Bermudagrass, egg weight, xanthophyll, yolk color, 1154
body weight, broiler breeder performance, lighting program, 48*
calcium, lead, quail-Japanese, 81*
bone, egg shell quality, feed intake, magnesium excess, 97*
brewers dried grains, fertility, hatchability, pheasant breeder, 138*
broiler breeder, feed restriction, sexual maturity, 1001
broiler breeder performance, fertility, hatchability, male-feeding system, 48*
floor type, hatchability, mortality, 1414
mortality, rabbit excreta, 39*
broody, egg quality, progesterone, turkey, 72*
cage area, corticosterone, dominance rank, layer performance, 399
calbindin, heat stress, layer, 52*
calcium source, limestone solubility, particle size, shell quality, 66*
chlorella, layer performance, 84*
control populations, egg weight, hatchability, quail-Japanese, 357
controlled feeding, energy utilization, 136*
dietary level, Haugh units, wheat middlings, 1329
dietary protein, layer performance, pullet dietary treatment, 62*
dietary pyruvic acid, fatty liver hemorrhagic syndrome,

feed intake, heat stress, 104*
disease challenge, *Pasteurella multocida*, turkey, 1807†
dry feed, heat stress, wet feed, 164*
dust-bathing activity, egg quality, fear-related response, quail-Japanese, 1363
egg mass, genetic parameters, selection index, shell strength, 42*
layer, lysine assay, soybean meal, 49*
egg quality, field beans, layer performance, vicine, 455
egg shell quality, layer, *Mycoplasma gallisepticum*, 5*
energy, protein, pullet growth, 62*
energy level, layer diet, temperature-hot weather, 108*
Ethacal® feed component, layer, phosphorus level, 86*
feathering, nervousness, pullet, rearing environment, 1400
feed conversion, fertility and hatchability, rabbit excreta, semen production, 168*
molt, photorefractoriness, thyroxine, turkey, 21*
propylthiouracil, pullet, sexual maturity, 107*
Egg production cycles, egg production models, nonlinear models, 213
Egg production models, egg production cycles, nonlinear models, 213
growth function models, 64*
Egg production rate, layer replacement, multiple hen cages, 848†
Egg products, *Campylobacter jejuni* contamination, egg-processing plant, 1431
dimethyl sulfide, odor, spoilage, uracil, 123*
Egg quality, age of bird, egg storage position, shell treatment, yolk position, 75*
broody, egg production, progesterone, turkey, 72*
dust-bathing activity, egg production, fear-related response, quail-Japanese, 1363
egg production, field beans, layer performance, vicine, 455
Egg shell, calcium reserve, structural unit, 77*
Egg shell calcification, 1,25-dihydroxycholecalciferol, egg shell quality, layer performance, 1322
Egg shell membranes, amino acid analysis, egg shell quality, shell-less egg, soft-shelled egg, 1203
Egg shell permeability, conductance, hatchability, turkey, 9*
hatchability, iodine, turkey egg, 67*
Egg shell quality, 1,25-dihydroxycholecalciferol, egg shell calcification, layer performance, 1322
1,25-dihydroxyvitamin D_3, layer performance, vitamin D, 166*
acid-base balance, dietary calcium, dietary chloride, sodium, 750
amino acid analysis, egg shell membranes, shell-less egg, soft-shelled egg, 1203
bone, egg production, feed intake, magnesium excess, 97*
bone strength, correlation, 132*
computerized data collection system, egg specific gravity, egg weight, 902
egg production, layer, *Mycoplasma gallisepticum*, 5*
egg specific gravity, sodium aluminosilicate, zeolite, 577
molt, pimpling, 205
Egg shell strength, molt-induced, phospholipids, shell gland, 427
Egg shell structure, abnormal egg shell structure, calcium reserve, 78*
palisade layer, 77*
Egg shell thickness, computer data collection, electronic gauge performance, 1105†
Egg shell weight, bone, calcium level, prelay pullet, 98*
Egg specific gravity, computerized data collection system, egg shell quality, egg weight, 902
egg shell quality, sodium aluminosilicate, zeolite, 577
Egg storage, cytogenetics, embryo abnormalities, heteroploid, karyotype, 96*
Egg storage position, age of bird, egg quality, shell treatment, yolk position, 75*
Egg turning, day of incubation, hatchability, high density plastic egg flat, malposition, 685
Egg type pullet, early protein restriction, growth restriction, layer performance, 54*
mash, pellets *vs.* mash, temperature, weight gain, 375
Egg weight, ahemeral lighting, egg production, lighting program, pullet, 176*
barley, feed utilization, layer performance, metabolizable energy, 1306
Bermudagrass, egg production, xanthophyll, yolk color, 1154
computerized data collection system, egg shell quality, egg specific gravity, 902
control populations, egg production, hatchability, quail-Japanese, 357
feed efficiency, Fermacto®, layer performance, methionine level, 842†
layer, molt-induced, zinc, 5*
shell calcium, shell thickness, shell weight, specific gravity, 159*
Egg weight loss, embryonic development, embryonic mortality, quail-Japanese, thiouracil, 28*
Egg white, chemical denaturation, egg white protein, gelation, 53*
Egg white protein, chemical denaturation, egg white, gelation, 53*
Egg yolk, age, layer, strain, 7*
choline supplementation, high performance liquid chromatography, methionine supplementation, phospholipids, 1717
food-poisoning bacteria, lecithovitellin reaction, salted yolk paste, *Staphylococcus aureus*, 261
physical manipulation, salt, shaking, texture, 1264
Egg yolk pigmentation, layer, xanthophyll, 126*
Egg yolk protein precursor, DNA modifications, quail-Japanese, vitellogenin gene, 70*
Eimeria, anticoccidial efficacy, battery cage trials, ionophorous antibiotic, narasin, 1043
coccidiosis, diclazuril, 115*
floor pen trials, ionophorous antibiotic, narasin, 1050
coccidiosis, immune response, recombinant coccidial antigens, 68*
foreign hosts, sporozoites, 49*
immune response, prokaryotic expression vector, recombinant DNA, 100*
Eimeria acervulina, antibody response, *B*-congenic chickens, p150 merozoite protein, recombinant coccidial antigens, 111*
apolipoprotein A-I, carotenoids, chick intestine, coccidiosis, 45*
chick growth, feed efficiency, lysine deficiency, 232
Eimeria tenella, antibody, broiler chick, coccidia, ELISA, 16*
arprinocid, broiler resistance, glycamide, nicarbazin, 164*
coccidiosis, lymphokines, 105*
recombinant DNA, vaccine, 72*
Ejaculate, semen production, spermatogenesis, turkey, 327
Elastin, aorta, *Bordetella avium*, trachea, turkey poult, 174*

Electric shock-experimental system, behavior, chick performance, health, stress, 1791†
Electric stimulation, accelerated rigor, minimum time process, nonaged tenderness, tenderness, 40*
adenosine triphosphate, pH, poultry carcass, tenderness, 106*
boning time, broiler breast fillets, post-mortem metabolism, tenderness, 150*
broiler breast meat, hot boning, poultry meat color, texture, 1536
broiler breast muscle, texture profile, voltage levels, 23*
broiler muscle, ion exchange, lactate dehydrogenase, 11*
prolactin, serotonin, turkey, 93*
Electric stunning, delayed glycolysis, muscle pH, protein solubility, turkey muscle, 1062
broiler breast meat, hot boning, meat quality, 674†
Electrocuting flies, black light devices, caged layer, flies, muscalure, 871
Electron microscopy, epididymis, gossypol acetic acid, quail-Japanese, testis, 961
Electronic gauge performance, computer data collection, egg shell thickness, 1105†
ELISA, acquired immunity, broiler, *Cryptosporidium baileyi*, immune response, 720
antisperm antibodies, immunohistochemistry, turkey fertility, 20*
antibody, broiler chick, coccidia, *Eimeria tenella*, 16*
bovine serum albumin, chick, dopamine, serotonin, 100*
broiler, monoclonal antibodies, *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, 116*
broiler carcass, feed, gene probe, *Salmonella* isolation, 921
broiler performance, cryptosporidiosis, *Cryptosporidium*, 730
coccidiosis, recombinant coccidial antigens, 103*
cooked chicken, *Listeria monocytogenes*, 157*
Embryo, bone development, egg injection, vitamin A, vitamin D_3, 58*
chlordecone, corticosterone, microwave radiation, quail-Japanese, 60*
hatching egg injection machine, incubation, Marek's disease, vaccination, 110*
Embryo abnormalities, cytogenetics, egg storage, heteroploid, karyotype, 96*
Embryogenesis, morphogenesis, pigeon, secondary plate development, 865
Embryonic cell transfer, chick embryo, chimera chick, 137*
Embryonic development, egg weight loss, embryonic mortality, quail-Japanese, thiouracil, 28*
Embryonic growth, feather growth, feather pigmentation, 56*
Embryonic mortality, abnormal eggs, cage *vs.* floor housing, fertility, turkey, 1455
egg weight loss, embryonic development, quail-Japanese, thiouracil, 28*
fertility, mineral level, 127*
high altitude, incubation temperature, oxygen tension, turkey egg hatchability, 956
Endocrine glands, light environment, light intensity, population density, turkey performance, 1518
Endogenous virus, avian leukosis virus, congenital transmission, sex-linkage, slow-feathering, 1668
slow-feathering, tumors, 1674
early feathering, genes, Z chromosome, 191
hemagglutination, R_2 alloantisera, 50*
Energy, amino acid(s), bioavailability, guinea fowl, Leghorn, 145*
body composition, protein, turkey breeder performance, 17*
broiler chick, carcass composition, compensatory growth, feed restriction, 61*
carcass composition, layer, metabolizable energy utilization, temperature, 136*
dietary protein, reproductive performance, triticale, turkey semen, 1619
egg production, protein, pullet growth, 62*
feed and water intake, feed efficiency, protein, quail-Japanese growth, 1111
pellets, protein level sequence, turkey tom performance, 170*
Energy consumption, air velocity, cage environment, feed intake, Leghorn cockerels, 62*
Energy estimation, carcass composition, lipid fraction, phospholipids, protein content, 413
Energy feeding regimen, abdominal fat, broiler chick performance, dressing percentage, intestinal measurement, 131*
Energy level, amino acid(s), broiler, sex separate rearing, 39*
egg production, layer diet, temperature-hot weather, 108*
layer diet, lighting program, prolapse, pullet diet, 108*
layer performance, pullet diet, 63*
protein level, pullet, self-select diets, sexual maturity, 153*
Energy metabolism, anorexia, feed intake, thiamin deficiency, turkey, 143*
Energy needs, feed intake, poultry production, 115*
Energy repletion, broiler chick, compensatory growth, enzyme activities, lipogenesis, 146*
Energy restriction, broiler crosses, compensatory growth, feed efficiency, 156*
Energy source, adipose tissue, broiler pullet, feed access, 766
Energy supplementation, confounding experimental results, growth assay, oil, sand, 169*
Energy use, broiler diet, feed efficiency, light source, weight gain, 47*
Energy utilization, controlled feeding, egg production, 136*
Enteropathogen, broiler chick, detection, *Salmonella typhimurium*, 52*
Environmental stress, layer performance, relative humidity, temperature, vitamin C, 67*
Environmental temperature, antibody production, immune response, selection lines, sheep red blood cells, 79*
ascites, broiler, roaster, 87*
bone meal source, broiler performance, broiler strain, phosphorus availability, 27*
broiler, digestive tract content, feed regimen, 64
Enzyme activities, broiler chick, compensatory growth, energy repletion, lipogenesis, 146*
Enzyme immunoassay, cell wall antigen, *Listeria monocytogenes*, 157*
Enzyme supplementation, barley, chick performance, viscosity, weight gain, 147*
Epidemiological evaluation, competitive exclusion, *Salmonella* control, 1026
Epidermal growth factor, gonadotrophins, granulosa cell, *in vitro*, steroid hormones, 814
Epididymis, electron microscopy, gossypol acetic acid, quail-Japanese, testis, 961
Epinephrine, catecholamines, dopamine, norepinephrine,

75*
Epiphyseal growth plate, broiler chick, growth hormone, tibial dyschondroplasia, 82*
Erythrocyte alloantigens, hemagglutination reactions, immunogenetic analysis, immunoglobulin G, mitogen response, 1210
Erythrocytes, 2,3-bisphosphoglycerate, inositol pentakisphosphate, oxygen affinity, phosphate, 100*
hemolysis, plasma, t-butyl hydroperoxide, uric acid, 1632
Escherichia coli, antibiotic resistance, food chain, kanamycin, *Salmonella*, 699
kanamycin, *Salmonella*, transconjugant, 689
attenuated vaccine, *Pasteurella multocida*, 113*
broiler performance, coliforms, Luprosil, 13*
cloning, *Mycoplasma gallisepiicum*, 26*
Congo Red negative, Congo Red positive, macrophages, 121*
Estradiol, broiler male, sudden death syndrome, 156†
ovarian growth, pregnant mare serum gonadotropin, prolactin, turkey, 319
Ethacal® feed component, body weight gain, broiler performance, temperature environment, 18*
broiler breeder performance, specific gravity, 35*
broiler chick, sodium availability, 15*
egg production, layer, phosphorus level, 86*
layer, sodium availability, 120*
Ethoxyquin, broiler performance, feed preservative, peroxide, rancidity, 1725
Euglycemic clamp technique, *B*-adrenergic agonist, chick, glucose, insulin, 470
Evaporative cooling, air misting, broiler, 8*
broiler, fan cooling, pad cooling, ventilation, 98*
Eviscerated carcass yield, abdominal fat, breast yield, broiler, feathering genotypes, 914
Eviscerated yield, abdominal fat, beta agonist, feeding regimen, quail-Japanese, 120*
abdominal fat pad, *am* gene, broiler, 102*
Evisceration, broiler processing, chilling, *Salmonella* control, scalding, 933
Excreta, Alimet®, broiler chick, DL-methionine, metabolism, 12*
methionine, 76*
Excreta collection, bioassay, glucose, true metabolizable energy, water, 1192
Exercise, atherosclerosis, cholesterol, quail-Japanese, 161*
broiler breast muscle, broiler drumsticks, broiler growth, muscle characteristics, 1557
Exercise patterns, breast muscle, broiler, muscle growth, thigh, 1708
Experimental surgery, double oviduct, fertility, oviposition, 1465
Expert decision support system, computer software, expert system technology, layer management, 152*
Expert system technology, computer software, expert decision support system, layer management, 152*
Extension poultry specialist role, economic development, 136*
extension program, poultry industry, poultry programs of the future, 883
Extension program, educational slides, egg grading and packing, egg-processing plant design, egg-processing slides, 54*
extension poultry specialist role, poultry industry, poultry programs of the future, 883
poultry programs of the future, poultry students, teaching poultry, 878
Exudate, marinade composition, salt soluble protein, tumbling, turkey meat, 162*
Fan cooling, broiler, evaporative cooling, pad cooling, ventilation, 98*
Fasted rooster, feather meal, metabolizable energy assays, variances, 1188
Fasting, broiler performance, heat stress, potassium chloride, weight gain, 36*
Fasting heat production, body size, body weight, 146*
Fasting periods, days without eggs, feather loss, layer performance, peaks of production, 67*
Fat, amino acid(s), protein, turkey, 4*
broiler performance, grain sorghum, metabolizable energy, tannin content, 80*
lecithin, phospholipids, shell gland, shell strength, 8*
Fat blends, broiler chick diet, poultry fat, yellow grease, 27*
Fat content, beef sausage, meat flavor, quality attributes, turkey sausage, 1075
Fat deposition, free fatty acids, glucagon, insulin, 69*
Fat free mass, bioelectrical impedance analysis, body composition, turkey breeder, 92*
Fat in diet, broiler breeder, fertility, hatchability, lighting program, 5*
Fat pad, amaranth, body weight, broiler male performance, processing yield, 743
Fat sources, fatty acid profile, lipid oxidation, sausage, 91*
Fatty acid composition, blended feed-grade fats, gas-liquid chromatography, *trans*/*cis* 18:1, 307
Fatty acid profile, fat sources, lipid oxidation, sausage, 91*
Fatty acids, biotin, broiler chick, phospholipids, 171*
chick, liver lipids, *trans*/18:1, 45*
broiler, carcass composition, carcass fat, 165*
corn, processing yield, sorghum, 130*
chick tissue, hydrogenated, *trans*/*cis* 18:1, 778
Fatty liver hemorrhagic syndrome, dietary pyruvic acid, egg production, feed intake, heat stress, 104*
glutathione supplementation, protein level, quail-Japanese, 159*
Fatty material injection, consumer preference, roasted turkey, turkey broth injection, 160*
Fawn, genetics, mutation, plumage color, quail-Japanese, 127*
Fear-related response, dust-bathing activity, egg production, egg quality, quail-Japanese, 1363
Feather color, photoperiod, quail-Japanese, testicular development, 1098†
Feather condition, diurnal rhythm, layer, lighting program, operant heat demand, 108*
Feather growth, embryonic growth, feather pigmentation, 56*
Feather loss, days without eggs, fasting periods, layer performance, peaks of production, 67*
Feather meal, abdominal fat, broiler, carcass measurement, lysine, 111*
abdominal fat deposition, broiler, finishing period, nonspecific nitrogen source, 300
fasted rooster, metabolizable energy assays, variances, 1188
Feather pigmentation, embryonic growth, feather growth, 56*
Feather release, broiler, microbiological contamination, picking feathers, spray scalding, 12*
Feather release forces, broiler, scalding temperature, scalding time, stunning *vs.* nonstunning, 1069
Feather removal, broiler crosses, feeding regimen, kosher processing, skin strength, 125*
Feathering, egg production, nervousness, pullet, rearing environment, 1400
Feathering genotypes, abdominal fat, breast yield, broiler, eviscerated carcass yield, 914

Feathers, chick, fiber optics, sex-linked coloration, 1680
Feed, broiler carcass, chemical preservative, Myco Curb®, *Salmonella*, 1225†
ELISA, gene probe, *Salmonella* isolation, 921
broiler chick performance, supplemental equipment, water, 124*
Feed access, adipose tissue, broiler pullet, energy source, 766
Feed additive, broiler diet, feed flavor enhancer efficacy, 22*
Feed and water intake, energy, feed efficiency, protein, quail-Japanese growth, 1111
Feed and water withdrawal, gastrointestinal tract, moisture content, shear strength, water withdrawal, 845†
Feed consumption, age, beak trimming, guinea fowl, mortality, 141†
beak trimming, behavior, pullet, weight gain, 109*
duodenum, layer, molt-induced, zinc, 58*
Feed conversion, abdominal fat, body weight, broiler sire strain, heritability, 1
selection, 10
broiler, growth rate, light intensity, 73*
broiler sire strain, slaughter yield, weight gain, 16
amino acid(s), body weight, broiler male chick, threonine requirement, 108
antibiotic, broiler performance, weight gain, 70*
body weight, broiler, wheat screenings, 615
broiler chick, soybean meal, tibial dyschondroplasia, 21*
leg abnormalities, niacin requirement, turkey poult, 760
egg production, fertility and hatchability, rabbit excreta, semen production, 168*
layer performance, phase feeding, protein dietary regimens, 1447
Feed deprivation, broiler chick, delayed hypersensitivity, feed intake-overconsumption, immunocompetence, 626
Feed efficiency, abdominal fat, age, broiler chick, feed restriction, 384
age and weight-constant efficiency, broiler growth, feed intake, weight correction, 565
age constant, body weight, heritability, weight constant, 1497
air sac lesions, broiler, fructooligosaccharides, mortality, 46*
Alimet®, brown egg layer, egg mass, methionine, 16*
attitudes of integrators, biosecurity, broiler industry survey, growth regulators, 50*
beak trimming, broiler chick performance, heterophil:lymphocyte ratio, stress, 166*
body weight, dietary fat, protein, turkey male, 1281
feed intake, protein restriction, turkey male performance, 130*
bone calcium, chick, coccidiosis, zeolite-A, 172*
breast blister, broiler performance, floor type, 48*
broiler, compensatory growth, feed restriction, 118*
feed restriction, growth rate, sudden death syndrome, 1102†
pelleting, sunflower meal, 175*
broiler crosses, compensatory growth, energy restriction, 156*
broiler diet, energy use, light source, weight gain, 47*
broiler male, floor pen reared, fructooligosaccharides, 1*
carcass composition, carcass yield, corn gluten feed, turkey performance, 585
carcass fat, dietary fat, growth rate, turkey, 420
chick growth, *Eimeria acervulina*, lysine deficiency, 232
egg weight, Fermacto®, layer performance, methionine level, 842†
energy, feed and water intake, protein, quail-Japanese growth, 1111
Feed flavor enhancer efficacy, broiler diet, feed additive, 22*
Feed form, floor space, lighting environment, turkey management, 127*
Feed inhibition, acetic and propionic acids, broiler chick, feed intake, 139*
Feed intake, acetic and propionic acids, broiler chick, feed inhibition, 139*
age and weight-constant efficiency, broiler growth, feed efficiency, weight correction, 565
air velocity, cage environment, energy consumption, Leghorn cockerels, 62*
animal fat, body weight, broiler chick performance, monensin, 452
anorexia, energy metabolism, thiamin deficiency, turkey, 143*
B-endorphin, body temperature, water intake, 25*
beak trimming, body weight, broiler roaster growth, pellets *vs.* mash, 1514
body fat, chick, dietary energy, meal feeding, 140*
body weight, feed efficiency, protein restriction, turkey male performance, 130*
bone, egg production, egg shell quality, magnesium excess, 97*
broiler, carbonated water, heat stress, mortality, 30*
force feeding, plasma free fatty acids, plasma ketones, 30*
heat stress, water intake, 38*
plasma free arachidonic acid, 144*
broiler chick growth, glycine, glycolic acid, glyoxylic acid, 1469
broiler performance, cyclic watering, water intake, 378
calcium intake, layer, self-select diets, 24*
catecholamines, chicken, peptides, turkey, 74*
chick, hypothalamus, neuropeptide, 107*
dietary pyruvic acid, egg production, fatty liver hemorrhagic syndrome, heat stress, 104*
energy needs, poultry production, 115*
Feed intake principles, care of poultry, management, 90*
Feed intake-overconsumption, broiler chick, delayed hypersensitivity, feed deprivation, immunocompetence, 626
Feed preservative, broiler performance, ethoxyquin, peroxide, rancidity, 1725
Feed regimen, broiler, digestive tract content, environmental temperature, 64
Feed restriction, abdominal fat, age, broiler chick, feed efficiency, 384
broiler performance, glycolic acid, growth, 28*
age, broiler, heat stress, mortality, 2*
aggressive behavior, broiler breeder male, tryptophan, 155*
body composition, broiler breeder, semen production, 34*
body weight, broiler breeder pullet, feeder space, 996
plasma corticosterone, plasma glucose, 809
molt, temperature, 137*
broiler, compensatory growth, feed efficiency, 118*
growth hormone, 118*
feed efficiency, growth rate, sudden death syndrome, 1102†
heat stress, mortality, 226
broiler breeder, corticosterone, stress, tryptophan, 119*
egg production, sexual maturity, 1001

intestinal adaptation, 49*
broiler chick, carcass composition, compensatory growth, energy, 61*
broiler performance, lighting program, skeletal diseases, 68*
carcass composition, compensatory growth, turkey male, 1407
corticosterone, quail-Japanese, testosterone, 121*
Feed trough partitions, behavior, cage shape, layer performance, 1348†
Feed utilization, barley, egg weight, layer performance, metabolizable energy, 1306
Feeder space, body weight, broiler breeder pullet, feed restriction, 996
Feeder type, cage type, hen density, layer performance, 102*
Feeding regimen, abdominal fat, beta agonist, eviscerated yield, quail-Japanese, 120*
body composition, broiler breeder male, semen characteristics, 835
broiler breeder, plasma calcium, plasma phosphorus, 56*
broiler crosses, feather removal, kosher processing, skin strength, 125*
Feeding time, broiler breeder, hatchability, shell quality, slow *vs.* fast feathering, 538
Leghorn male, meal feeding, plasma, uric acid, 345†
Feedstuffs, ash, bomb calorimeter, proximate analysis, 513†
Fermacto®, egg weight, feed efficiency, layer performance, methionine level, 842†
Fermentation, carbon sources, coliforms, manure-cage layer, *Salmonella*, 74*
Fertility, abnormal eggs, cage *vs.* floor housing, embryonic mortality, turkey, 1455
artificial insemination, semen extender, spermatozoa-damaged, turkey semen, 1483†
biotin level, hatchability, turkey egg, 30*
brewers dried grains, egg production, hatchability, pheasant breeder, 138*
broiler breeder, fat in diet, hatchability, lighting program, 5*
broiler breeder egg, hatchability, zeolite-A, 19*
broiler breeder performance, egg production, hatchability, male-feeding system, 48*
carboxymethyl cellulose, semen extender, semen viscosity, 1218
corn-soybean meal diet, hatchability, niacin supplementation, turkey breeder performance, 336†
double oviduct, experimental surgery, oviposition, 1465
embryonic mortality, mineral level, 127*
freezing, semen, spermatozoal organelles, ultrastructure, 819
frozen semen, storage container, 333†
hatchability, immunogenetics, inbreeding, sublines, 859
sanitizers, turkey egg, 144*
mineral level in eggs, mineral level in hen blood, 163*
naked neck, reproductive traits, temperature, 176*
poly-L-glutamic acid, 82*
Fertility and hatchability, egg production, feed conversion, rabbit excreta, semen production, 168*
Fertilization, acrosome reaction, cock spermatozoa, *in vitro* incubation, perivitelline layer, 1770
FFA and 4-H, poultry majors, poultry programs of the future, student recruitment, youth programs, 886
Fiber optics, chick, feathers, sex-linked coloration, 1680
Fiber orientation, broiler breast tissue, salt concentration, 122*
Fiber type, broiler muscle, chilling, rigor mortis, 150*
Field beans, egg production, egg quality, layer performance, vicine, 455
Field trials, anticoccidial efficacy, monensin, narasin, 1058
Filtration, broiler, chiller water, diatomaceous earth, processing plant, 155*
Financial attractiveness, computer spreadsheet templates, contract, turkey growout, 142*
Fingerprinting, infectious bronchitis virus, protein, ribonucleic acid, 61*
Finishing period, abdominal fat deposition, broiler, feather meal, nonspecific nitrogen source, 300
Fitting guidelines, 4-H, duck, showmanship guidelines, turkey, 167*
Flies, black light devices, caged layer, electrocuting flies, muscalure, 871
Floor pen reared, broiler male, feed efficiency, fructooligosaccharides, 1*
Floor pen trials, anticoccidial efficacy, *Eimeria*, ionophorous antibiotic, narasin, 1050
Floor space, feed form, lighting environment, turkey management, 127*
Floor type, breast blister, broiler performance, feed efficiency, 48*
broiler breeder performance, egg production, hatchability, mortality, 1414
litter condition, turkey performance, 66*
Florescent, broiler growth performance, incandescent, light source, sodium vapor, 43
Flow cytometer, *Mycoplasma gallisepticum* identification, *Mycoplasma synoviae* identification, 25*
Flow cytometric analysis, *B*-congenic chickens, Ia antigens, lymphocyte response, monoclonal antibody, 1525
Flow rate, broiler chick, nipple waterer, 137*
Flow volume measurement, poultry-processing plant, water conservation, 119*
Fluoride, bone ash, bone strength, bone structure, broiler performance, 950
Fluorometry, broiler breeder, divergent selection, sperm viability, 55*
Follicle-stimulating hormone, granulosa cell, inhibin activity, luteinizing hormone, 1625
radioreceptor assay, turkey, 95*
Food chain, antibiotic resistance, *Escherichia coli*, kanamycin, *Salmonella*, 699
Food-poisoning bacteria, egg yolk, lecithovitellin reaction, salted yolk paste, *Staphylococcus aureus*, 261
Food Safety and Inspection Service, poultry inspection, *Salmonella* contamination in poultry, *Salmonella* control, 944
Foodborne disease costs, economic costs, risk assessment, *Salmonella* control, salmonellosis, 936
Foot temperature, conductive heat loss, heat stress, layer, thermal perch, 142*
Force feeding, broiler, feed intake, plasma free fatty acids, plasma ketones, 30*
Force resting, iron, layer performance, zinc, 3*
Foreign hosts, *Eimeria*, sporozoites, 49*
Formaldehyde, blood chemistry, layer, pellet binder, urea, 37*
hatching egg sanitizer, hydrogen peroxide, ozonated water, quaternary ammonium compound, 155*
Formic acid, broiler diet, calcium formate, fumaric acid, organic acids, 1178
Fowl cholera, dietary protein restriction, humoral immunity, *Pasteurella multocida*, 135*
KSCN extract, *Pasteurella multocida*, turkey, vaccination, 26*

Franubet cultivar, barley, beta-glucanase, beta-glucans, broiler chick growth, 1573
Free fatty acids, fat deposition, glucagon, insulin, 69*
Freezing, fertility, semen, spermatozoal organelles, ultrastructure, 819
Frozen semen, fertility, storage container, 333†
Fructooligosaccharides, air sac lesions, broiler, feed efficiency, mortality, 46*
broiler male, feed efficiency, floor pen reared, 1*
Frying conditions, chicken patties, lipid content, moisture content, shortening quality, 9*
Full-fat canola seed, Avizyme®, barley, broiler diet, layer diet, 156*
canola seed, pelleting, seed-processing methods, turkey, 1731
cholesterol, egg, omega-3 fatty acids, serum, 129*
Fumaric acid, broiler diet, calcium formate, formic acid, organic acids, 1178
Functional properties, composition, mechanically deboned meat, washing, 87*
Functional properties of myofibrils, lipid oxidation, myofibrils, storage time, water activity, 128*
Fusarochromanone, copper, dyschondroplasia, immunity, mycotoxin, 174*
Futher processing, microbiological growth support, nutrient availability, particle size, turkey meat product form, 1269
Gas-liquid chromatography, blended feed-grade fats, fatty acid composition, *trans/cis* 18:1, 307
Gastric mucosal cells, chick, cyclic adenosine-3′,5′-monophosphate, gizzard erosion, gizzerosine, 1290
Gastrointestinal tract, feed and water withdrawal, moisture content, shear strength, water withdrawal, 845†
Gel-strength, mechanically deboned meat, poultry meat color, surimi-wash process, 91*
Gelation, chemical denaturation, egg white, egg white protein, 53*
Gene expression, 5-azacytidine, DNA methylation, prolactin, turkey, 1794†
Gene interactions, black-tailed red pattern, dark brown gene, plumage color, Villafranquina chickens, 351
genetics, muscle growth, muscular dystrophy, 367
Gene mapping, *B*-actin gene, *in situ* hybridization, metaphase chromosome, 154*
Gene probe, broiler carcass, ELISA, feed, *Salmonella* isolation, 921
Gene transfer, chicken embryo, retrovirus vector, tissue-specific expression, 34*
chloramphenicol acetyltransferase, retrovirus vector, tissue-specific expression, 156*
retrovirus vector, selectable marker, 42*
Genes, avian leukosis virus, early feathering, endogenous virus, Z chromosome, 191
Genetic analysis, *Staphylococcus aureus*, wattle reaction, 70*
Genetic associations, defective eggs, oviposition rate, shell quality, shell-less egg, 123*
Genetic correlation, abdominal fat, lipase, lipoprotein, protein, 183
body weight, carcass composition, heritability, turkey, 1388
growth hormone, heritability, plasma, 1781†
heritability, skeletal measurments, turkey, 1378
selection, sexual dimorphism, turkey, 166*
Genetic lines, growth hormone, turkey growth rate, 826
immunity, intracerebral inoculation, neurological behavior, Rous sarcoma virus, 1123
Genetic merit, adult selection, juvenile selection, layer performance, selection response, 1655
Genetic parameters, egg mass, egg production, selection index, shell strength, 42*
Genetic selection, atherosclerosis, cholesterol, quail-Japanese, 51*
body weight, correlation, heat stress, 83*
growth, heat stress, resistance, 83*
Genetics, Fawn, mutation, plumage color, quail-Japanese, 127*
gene interactions, muscle growth, muscular dystrophy, 367
guinea fowl, *Numida melagris*, plumage color, 158*
Pavo cristatus, peafowl, plumage color mutation, sex-linked gene, 158*
Gentian violet, antifungal activity, heat, moisture, particle size, 58
broiler, chlortetracycline, T-2 toxin, 99*
Gizzard erosion, broiler chick growth, gizzerosine, mortality, 1580
chick, cyclic adenosine-3′,5′-monophosphate, gastric mucosal cells, gizzerosine, 1290
Gizzerosine, broiler chick growth, gizzard erosion, mortality, 1580
chick, cyclic adenosine-3′,5′-monophosphate, gastric mucosal cells, gizzard erosion, 1290
Glomerular size, kidney anatomy, pullet strain, watering device, 31*
Glucagon, fat deposition, free fatty acids, insulin, 69*
insulin, lame *vs.* normal, leg abnormalities, turkey, 97*
Glucose, *B*-adrenergic agonist, chick, euglycemic clamp technique, insulin, 470
bioassay, excreta collection, true metabolizable energy, water, 1192
glycogen, intestine, turkey poult, yolk sac, 493
Glucose supplement, antibiotic treatment, hatching metabolism, turkey embryo, 463
Glutathione peroxidase, broiler chick, glutathione reductase, riboflavin, selenium, 149*
roxarsone, selenium, 41*
Glutathione reductase, broiler chick, glutathione peroxidase, riboflavin, selenium, 149*
Glutathione supplementation, fatty liver hemorrhagic syndrome, protein level, quail-Japanese, 159*
Glycamide, arprinocid, broiler resistance, *Eimeria tenella*, nicarbazin, 164*
Glycerol, broiler breeder evaluation, semen, tissue culture, 23*
cell culture, spermatozoa-chicken, 135
Glycerol trioleate, pancreatic lipase, turkey, 85*
Glycine, broiler chick growth, feed intake, glycolic acid, glyoxylic acid, 1469
Glycogen, glucose, intestine, turkey poult, yolk sac, 493
Glycogen content, broiler breast muscle, hot boning, lactate content, proximate analysis, 1702
Glycogenolysis, buffers, hormones, turkey liver explants, 147*
Glycol solution, broiler, chilling time, packaged carcasses, temperature, 154*
Glycolic acid, abdominal fat, broiler performance, feed restriction, growth, 28*
broiler chick growth, feed intake, glycine, glyoxylic acid, 1469
Glycoprotein, egg, membrane layers, ovomucin, vitelline membrane, 476
Glycoprotein hormone, hybridoma, luteinizing hormone, monoclonal antibodies, turkey, 117*
Glyoxylic acid, broiler chick growth, feed intake, glycine, glycolic acid, 1469

Goals and strategy, Coturnix International, educational program, youth development, 167*
Gonadotroph, D-Trp-6-LHRH, layer, molt, reproductive performance, 38*
Gonadotrophins, epidermal growth factor, granulosa cell, *in vitro*, steroid hormones, 814
Goose embryo, 3,3′,4,4′-tetrachlorobiphenyl, duck embryo, herring gull embryo, teratogenicity, 52
Gossypol acetic acid, electron microscopy, epididymis, quail-Japanese, testis, 961
Graft *vs.* host reaction, adrenocorticotropin, ascorbic acid, cell-mediated immunity, splenomegaly, 313
Grain sorghum, broiler performance, fat, metabolizable energy, tannin content, 80*
Granulosa cell, epidermal growth factor, gonadotrophins, *in vitro*, steroid hormones, 814
follicle-stimulating hormone, inhibin activity, luteinizing hormone, 1625
growth factors, layer, plasminogen activator, 165*
Grass-clipping length, carotene, drying method, turf Bermudagrass, xanthophyll, 1644†
Grit, barley, corn texture, pellets, turkey performance, 101*
Grower organization, industry regulation, nuisance abatement, peer pressure, 169*
Growout process, broiler performance, productivity, quality, 4*
Growth, abdominal fat, broiler performance, feed restriction, glycolic acid, 28*
antibiotic, bacitracin, body weight, guinea fowl, 145†
color-filtered light, photoperiod, turkey performance, 1505
genetic selection, heat stress, resistance, 83*
muscle development, turkey genotypes, 22*
myopathies, turkey muscle, 173*
nutritional requirements, quail-Japanese, selection, 22*
Growth assay, arginine utilization, chick, lysine utilization, monensin, 341†
confounding experimental results, energy supplementation, oil, sand, 169*
Growth curve, multiphasic analysis, weight gain, 33
Growth factors, granulosa cell, layer, plasminogen activator, 165*
Growth function models, egg production models, 64*
Growth hormone, adenohypophysis, age, chick, somatotrophs, 487
age, metabolic clearance rate, protein restriction, secretion rate, 120
antibody production, leukocyte distribution, lymphoid organs, T_3,T_4 doses, 92*
B-cells, natural killer cells, T-cells, T_3,T_4 doses, 93*
body weight, genetic correlation, heritability, plasma, 1781†
broiler, compensatory growth, feed restriction, 118*
insulin-like growth factor-1, plasma hormone level, 117*
broiler chick, epiphyseal growth plate, tibial dyschondroplasia, 82*
broiler pullet, carcass composition, pulsatile, insulin, T_3,T_4, 168*
chick embryo, cholesterol, insulin-like growth factor-1, very low density lipoprotein, 71*
cloning, nucleotide sequence, 107*
genetic lines, turkey growth rate, 826
Growth performance, anticoccidial drugs, broiler, coccidial lesion scores, litter moisture, 126*
barley, beta-glucanase, broiler, 7*
Growth rate, abdominal fat, broiler, feed conversion, light intensity, 73*
amino acid(s), lysine, methionine, turkey tom performance, 86*
broiler, feed efficiency, feed restriction, sudden death syndrome, 1102†
carcass fat, dietary fat, feed efficiency, turkey, 420
heritability, protein and energy, quail-Japanese, selection differentials, 519
Growth regulators, attitudes of integrators, biosecurity, broiler industry survey, feed efficiency, 50*
Growth restriction, early protein restriction, egg type pullet, layer performance, 54*
Guinea fowl, age, beak trimming, feed consumption, mortality, 141†
amino acid(s), bioavailability, energy, Leghorn, 145*
antibiotic, bacitracin, body weight, growth, 145†
carcass yield, meat characteristics, 82*
genetics, *Numida melagris*, plumage color, 158*
ovarian follicle growth, pigeon, 1783†
Gypsum, acid-base balance, layer diet, pulverized and particle size limestone, shell quality, 20*
Halofuginone, anticoccidial drugs, coccidiosis, nicarbazin, polyether ionophores, 31*
broiler performance, maduramicin, monensin, nicarbazin, 32*
Hardness, chewiness, cooking method, deboning time, *pectoralis* muscle, 112*
Hatch time, house fly, larvicide, manure, thiourea, 407
Hatch weight, dry bulb temperature, wet bulb temperature, 37*
Hatchability, age at sexual maturity, dietary protein, dwarf broiler breeder performance, reproductive performance, 219
biotin level, fertility, turkey egg, 30*
brewers dried grains, egg production, fertility, pheasant breeder, 138*
broiler breeder, fat in diet, fertility, lighting program, 5*
feeding time, shell quality, slow *vs.* fast feathering, 538
broiler breeder egg, fertility, zeolite-A, 19*
broiler breeder performance, egg production, fertility, male-feeding system, 48*
floor type, mortality, 1414
conductance, egg shell permeability, turkey, 9*
control populations, egg production, egg weight, quail-Japanese, 357
corn-soybean meal diet, fertility, niacin supplementation, turkey breeder performance, 336†
day of incubation, egg turning, high density plastic egg flat, malposition, 685
egg shell permeability, iodine, turkey egg, 67*
fertility, immunogenetics, inbreeding, sublines, 859
sanitizers, turkey egg, 144*
Hatching egg injection machine, embryo, incubation, Marek's disease, vaccination, 110*
Hatching egg sanitizer, formaldehyde, hydrogen peroxide, ozonated water, quaternary ammonium compound, 155*
hydrogen peroxide, ozonated water, quaternary ammonium compound, 57*
Hatching metabolism, antibiotic treatment, glucose supplement, turkey embryo, 463
Hatching time, bone mineralization, broiler chick, 59*
broiler chick hatch weight, dry bulb temperature, wet bulb temperature, 163*
Haugh units, dietary level, egg production, wheat middlings, 1329
Health, behavior, caging system-experimental, chick performance, multiple stressors, 1787†

chick performance, electric shock-experimental system, stress, 1791†
Heat, antifungal activity, gentian violet, moisture, particle size, 58
Heat resistance, broiler, computer controlled environment, heat stress, 12*
Heat stress, acid-base balance, broiler, nicarbazin, 3*
acid-base status, blood, layer, thermoregulation, 113*
age, broiler, feed restriction, mortality, 2*
antibody production, immune response, selection lines, sheep red blood cells, 79*
arginine vasotocin, broiler, mesotocin, surgery stress, 171*
ascorbic acid, aspirin, broiler male chick growth, 1183
body temperature, naked neck *vs.* normal, 818
body weight, correlation, genetic selection, 83*
naked neck, 13*
broiler, carbonated water, feed intake, mortality, 30*
computer-controlled environment, heat resistance, 12*
feed intake, water intake, 38*
feed restriction, mortality, 226
broiler male performance, Maxiban®, mortality, nicarbazin, 17*
broiler meat, cooking yields, moisture pick-up, 27*
broiler performance, coccidiosis, narasin, nicarbazin, 248
fasting, potassium chloride, weight gain, 36*
calbindin, egg production, layer, 52*
conductive heat loss, foot temperature, layer, thermal perch, 142*
metabolic rate, panting, roost temperature, 124*
dietary pyruvic acid, egg production, fatty liver hemorrhagic syndrome, feed intake, 104*
dry feed, egg production, wet feed, 164*
genetic selection, growth, resistance, 83*
hypothalamus, layer performance, luteinizing hormone, luteinizing hormone releasing hormone, 79*
layer, *Mycoplasma gallisepticum*, 34*
Heat stress resistance, broiler, 14*
Hemagglutination, avian leukosis virus, endogenous virus, R_2 alloantisera, 50*
Hemagglutination reactions, erythrocyte alloantigens, immunogenetic analysis, immunoglobulin G, mitogen response, 1210
Heme-complex, cytochrome c, ligand, myoglobin, poultry meat color, 44*
hemoglobin, myoglobin, pH, poultry meat color, 44*
nicotinamide, poultry meat color, pyridine, turkey breast meat, 44*
Hemoglobin, heme-complex, myoglobin, pH, poultry meat color, 44*
Hemolysis, erythrocytes, plasma, t-butyl hydroperoxide, uric acid, 1632
Hen density, cage type, feeder type, layer performance, 102*
Hen plasma, steroid hormones, 167*
Heparin, nonesterified fatty acids, turkey, very low density lipoprotein, 51*
Hepatic tocopherol concentration, dietary fat, liver, turkey, vitamin E, 1814†
Heritability, abdominal fat, body weight, broiler sire strain, feed conversion, 1
age constant, body weight, feed efficiency, weight constant, 1497
body weight, carcass composition, genetic correlation, turkey, 1388
genetic correlation, growth hormone, plasma, 1781†
skeletal measurments, turkey, 1378
corticosterone, quail-Japanese, stress, 25
growth rate, protein and energy, quail-Japanese, selection differentials, 519
Herring gull embryo, 3,3′,4,4′-tetrachlorobiphenyl, duck embryo, goose embryo, teratogenicity, 52
Herring meal, cholesterol, layer performance, omega-3 fatty acids, 99*
Heterophil:lymphocyte ratio, beak trimming, broiler chick performance, feed efficiency, stress, 166*
Heterophils, antibody, *Brucella abortus*, humoral immunity, male chickens, 1775†
Heteroploid, cytogenetics, egg storage, embryo abnormalities, karyotype, 96*
High altitude, embryonic mortality, incubation temperature, oxygen tension, turkey egg hatchability, 956
High density plastic egg flat, day of incubation, egg turning, hatchability, malposition, 685
High energy electrons, chicken meat, irradiation pasteurization, *Salmonella*, 165*
High performance liquid chromatography, choline supplementation, egg yolk, methionine supplementation, phospholipids, 1717
Hindgut, Alimet®, broiler chick, DL-methionine absorption, 1314
Histidine, chick, copper, zinc, 47*
Histochemistry, muscle fiber type, skeletal muscle, 973
Histocompatibility complex, antitumor response, *B* complex genotype, Rous sarcoma virus, 164*
antibody response, immunocompetence, phagocytosis, T cell-mediated response, 989
Hormones, buffers, glycogenolysis, turkey liver explants, 147*
Host immunity, bovine serum albumin injection, northern fowl mite, rocket electrophoresis, two-dimensional immunoelectrophoresis, 1015
Hot boning, brine solution *vs.* water, broiler breast fillets, chilling temperature, tenderness, 1545
broiler breast meat, electric stimulation, poultry meat color, texture, 1536
electrical stunning, meat quality, 674†
broiler breast muscle, glycogen content, lactate content, proximate analysis, 1702
House fly, hatch time, larvicide, manure, thiourea, 407
Humoral immunity, antibody, *Brucella abortus*, heterophils, male chickens, 1775†
dietary protein restriction, fowl cholera, *Pasteurella multocida*, 135*
Hybridoma, glycoprotein hormone, luteinizing hormone, monoclonal antibodies, turkey, 117*
Hydrated sodium calcium aluminosilicate, aflatoxin, body weight, chick diet, 106*
Hydrogen peroxide, BOD removal, COD removal, poultry-processing water, wastewater treatment, 24*
formaldehyde, hatching egg sanitizer, ozonated water, quaternary ammonium compound, 155*
hatching egg sanitizer, ozonated water, quaternary ammonium compound, 57*
Hydrogenated, chick tissue, fatty acids, *trans*/*cis* 18:1, 778
Hydroxyechinenone, canthaxanthin, chick, isozeaxanthin, metabolism, 787
Hyperplastic response, abdominal fat, adipocyte hyperplasia, adipocyte hypertrophy, caloric overconsumption, 800
Hypersensitivity, broiler chick, immune response, soy protein, 104*
Hyperthermia, broiler, hypocalcemia, lactate, respiratory alklalosis, 132*

Hypocalcemia, broiler, hyperthermia, lactate, respiratory alklalosis, 132*
Hypophysectomy, 17α-hydroxylase, ovarian follicle, quail-Japanese, theca cell, 1636
Hypothalamus, biogenic amines, duodenum, mortality, turkey poult, 13*
chick, feed intake, neuropeptide, 107*
heat stress, layer performance, luteinizing hormone, luteinizing hormone releasing hormone, 79*
lesions, obesity, sexual dysfunction, 157*
photostimulation, prolactin, turkey, 138*
Ia antigens, *B*-congenic chickens, flow cytometric analysis, lymphocyte response, monoclonal antibody, 1525
Immune response, acquired immunity, broiler, *Cryptosporidium baileyi*, ELISA, 720
antibody, layer, northern fowl mite, Western blot analysis, 549
antibody production, environmental temperature, selection lines, sheep red blood cells, 79*
heat stress, selection lines, sheep red blood cells, 79*
light regimen, thymidine uptake, white blood cell, 114*
biological activity, interleukin-1, purification, 104*
broiler chick, hypersensitivity, soy protein, 104*
broiler egg, incubation, microwave radiation, 57*
bursal cell transfer, dopamine, plaque-forming cell transfer, sheep red blood cells, 88*
chick, zinc deficiency, 139*
coccidiosis, *Eimeria*, recombinant coccidial antigens, 68*
Eimeria, prokaryotic expression vector, recombinant DNA, 100*
Immunity, copper, dyschondroplasia, fusarochromanone, mycotoxin, 174*
genetic lines, intracerebral inoculation, neurological behavior, Rous sarcoma virus, 1123
infectious bronchitis virus, virus isolation, 174*
Immunization, antibody level, coccidiosis, protection antigen, 147*
Immunoblot, infectious bronchitis virus, 158*
Immunocompetence, antibody response, histocompatibility complex, phagocytosis, T cell-mediated response, 989
broiler chick, delayed hypersensitivity, feed deprivation, feed intake-overconsumption, 626
Immunogenetic analysis, erythrocyte alloantigens, hemagglutination reactions, immunoglobulin G, mitogen response, 1210
Immunogenetics, fertility, hatchability, inbreeding, sublines, 859
Immunoglobulin G, erythrocyte alloantigens, hemagglutination reactions, immunogenetic analysis, mitogen response, 1210
Immunohistochemistry, anti-sperm antibodies, ELISA, turkey fertility, 20*
Immunosuppression, chick, cyclosporine, salmonellosis, 176*
***In situ* hybridization**, *B*-actin gene, gene mapping, metaphase chromosome, 154*
In vitro, adrenal, adrenocorticotropin, bursectomy, chick, 83*
carrageenan, mitogenesis, peripheral blood lymphocytes, phytohemagglutinin-M, 33*
epidermal growth factor, gonadotrophins, granulosa cell, steroid hormones, 814
***In vitro* digestion**, amino acid(s), bioavailability, plant feedstuffs, true amino acid availability, 78
***In vitro* incubation**, acrosome reaction, cock spermatozoa, fertilization, perivitelline layer, 1770
Inbreeding, fertility, hatchability, immunogenetics, sublines, 859
Incandescent, broiler growth performance, fluorescent, light source, sodium vapor, 43
Incubation, broiler egg, immune response, microwave radiation, 57*
embryo, hatching egg injection machine, Marek's disease, vaccination, 110*
Incubation temperature, embryonic mortality, high altitude, oxygen tension, turkey egg hatchability, 956
Industry regulation, grower organization, nuisance abatement, peer pressure, 169*
Infectious bronchitis virus, antiviral, infectious bursal disease virus, 40*, 173*
chick, recombinant DNA, viral diagnosis, 69*
dietary calcium, kidney function, pullet chick, urolithiasis, 1250
fingerprinting, protein, ribonucleic acid, 61*
immunity, virus isolation, 174*
immunoblot, 158*
pathogenesis, tissue tropism, 60*
Infectious bursal disease virus, antiviral, infectious bronchitis virus, 40*, 173*
Inhibin activity, follicle-stimulating hormone, granulosa cell, luteinizing hormone, 1625
Inositol pentakisphosphate, 2,3-bisphosphoglycerate, erythrocytes, oxygen affinity, phosphate, 100*
Insulin, *B*-adrenergic agonist, chick, euglycemic clamp technique, glucose, 470
broiler pullet, growth hormone, pulsatile, T_3,T_4, 168*
chick, organ culture, pancreas, 101*
fat deposition, free fatty acids, glucagon, 69*
glucagon, lame *vs.* normal, leg abnormalities, turkey, 97*
Insulin-like growth factor-1, broiler, growth hormone, plasma hormone level, 117*
chick embryo, cholesterol, growth hormone, very low density lipoprotein, 71*
Interleukin-1, biological activity, immune response, purification, 104*
Internal quality, egg, storage condition, 133*
Interregional competition, broiler grower, profitability, simulation model, 89*
Intestinal adaptation, broiler breeder, feed restriction, 49*
Intestinal disaccharidases, age, diet, turkey poult, 153*
Intestinal measurement, abdominal fat, broiler chick performance, dressing percentage, energy feeding regimen, 131*
Intestinal spillage, bacteria, broiler carcass speck count, corn, milo, 88*
Intestinal uptake, broiler chick, methionine hydroxy analog, rat, 56*
Intestine, bile acids, broiler chick, liver, SEHCAT, 17*
broiler, location, Meckel's diverticulum, yolk stalk, 677†
chick, mannose sensitivity, receptors, *Salmonella typhimurium*, 133*
glucose, glycogen, turkey poult, yolk sac, 493
Intracerebral inoculation, genetic lines, immunity, neurological behavior, Rous sarcoma virus, 1123
Iodine, egg shell permeability, hatchability, turkey egg, 67*
Iodine supplementation, abdominal fat, bacitracin MD, body weight, broiler, 161*
Ion balance, chloride, layer diet, shell quality, sodium, 39*
Ion exchange, broiler muscle, electric stimulation, lactate dehydrogenase, 11*
Ionophores, bone ash, broiler chick, magnesium, 59*
Ionophorous antibiotic, anticoccidial efficacy, battery cage

trials, *Eimeria*, narasin, 1043
Eimeria, floor pen trials, narasin, 1050
Iraq rations, broiler performance, corn substitution, local barley, 122*
Iraqi chickens, semen, *Turella* yeast, 46*
Iron, force resting, layer performance, zinc, 3*
Irradiation, poultry products, *Salmonella*, sensory quality, shelf life, 36*
sensory properties, sodium reduction, turkey frankfurters, 1797†
Irradiation pasteurization, chicken meat, high energy electrons, *Salmonella*, 165*
Isoleucine, broiler chick, leg abnormalities, leucine, valine, 85*
Isoproterenol, broiler performance, carcass composition, weight gain, 95*
Isozeaxanthin, canthaxanthin, chick, hydroxyechinenone, metabolism, 787
Jackbean urease enzyme, antibiotics, antiurease antibody, chick growth, passive immunity, 434
Japan, layer performance year round, open-sided houses, photoperiod, seasonal change, 1130
Japanese quail, (see quail-Japanese)
Judging contest, computerized contest scoring, scan sheet scoring, 140*
Juvenile selection, adult selection, genetic merit, layer performance, selection response, 1655
Kamaboko-like products, composition, mechanically deboned meat, poultry meat color, yield, 112*
Kanamycin, antibiotic resistance, *Escherichia coli*, food chain, *Salmonella*, 699
Salmonella, transconjugant, 689
Karyotype, cytogenetics, egg storage, embryo abnormalities, heteroploid, 96*
Kidney anatomy, glomerular size, pullet strain, watering device, 31*
Kidney function, citrinin, diuresis, manure moisture, ochratoxin A, 1034
dietary calcium, infectious bronchitis virus, pullet chick, urolithiasis, 1250
Kosher processing, broiler crosses, feather removal, feeding regimen, skin strength, 125*
KSCN extract, fowl cholera, *Pasteurella multocida*, turkey, vaccination, 26*
L-methionine, broiler chick, DL-methionine, DL-methionine hydroxy analog, urinary excretion, 14*
Lactate, broiler, hyperthermia, hypocalcemia, respiratory alkalosis, 132*
broiler breast meat, denervation, pH, R value, 117*
Lactate content, broiler breast muscle, glycogen content, hot boning, proximate analysis, 1702
Lactate dehydrogenase, broiler muscle, electric stimulation, ion exchange, 11*
Lactic acid, bactericide, poultry-processing water, *Salmonella*, 69*
Lame *vs.* normal, glucagon, insulin, leg abnormalities, turkey, 97*
Larvicide, hatch time, house fly, manure, thiourea, 407
Lasalocid, bacitracin, body weight, turkey, 99*
Lathyrism, butylated hydroxyanisole, cysteine, quail-Japanese chick, toxicity, 153†
Layer, acid-base status, blood, heat stress, thermoregulation, 113*
age, egg yolk, strain, 7*
methionine requirement, sulfur amino acid requirement, 170*
agonistic activity, cage density, dominance rank, egg production, 198
antibody, immune response, northern fowl mite, Western blot analysis, 549
antinutritive effects, cecectomy, rapeseed meal, 61*
biliary secretion, pancreatic polypeptide, pancreatic secretion, 126
blood chemistry, formaldehyde, pellet binder, urea, 37*
blood levels, calcium, phosphorus, shell quality, 68*
bone, magnesium excess, phosphorus, plasma, 18*
broiler, economics, naked oats pricing, turkey, 10*
calbindin, egg production, heat stress, 52*
calcitriol, shell strength, 65*
calcium, phosphorus absorption, phosphorus excretion, 440
calcium intake, feed intake, self-select diets, 24*
carcass composition, energy, metabolizable energy utilization, temperature, 136*
chemical composition, culls, mortality, 94*
chloride level, performance index, phosphorus level, potassium level, 131*
clearance kinetics, narasin, 8*
conductive heat loss, foot temperature, heat stress, thermal perch, 142*
D-Trp-5-LHRH, gonadotroph, molt, reproductive performance, 38*
diurnal rhythm, feather condition, lighting program, operant heat demand, 108*
duodenum, feed consumption, molt-induced, zinc, 58*
egg assay for riboflavin, nutritional status, riboflavin, 160*
egg mass, egg production, lysine assay, soybean meal, 49*
egg production, egg shell quality, *Mycoplasma gallisepticum*, 5*
Ethacal® feed component, phosphorus level, 86*
egg weight, molt-induced, zinc, 5*
egg yolk pigmentation, xanthophyll, 126*
Ethacal® feed component, sodium availability, 120*
granulosa cell, growth factors, plasminogen activator, 165*
heat stress, *Mycoplasma gallisepticum*, 34*
luteinizing hormone, ovarian follicle, steroidogenesis, 145*
metabolizable energy intake, prediction equation, 135*
Layer diet, acid-base balance, gypsum, pulverized and particle size limestone, shell quality, 20*
Avizyme, barley, broiler diet, full-fat canola seed, 156*
broiler diet, Mazidagi raw rock phosphate, phosphorus source, 154*
chemical composition, naked oats, weathering, 115*
chloride, ion balance, shell quality, sodium, 39*
egg production, energy level, temperature-hot weather, 108*
energy level, lighting program, prolapse, pullet diet, 108*
Layer digestive system, calcium source, limestone, particle size, solubility, 142*
Layer management, computer software, expert decision support system, expert system technology, 152*
Layer performance, (see egg production also)
1,25-dihydroxycholecalciferol, egg shell calcification, egg shell quality, 1322
1,25-dihydroxyvitamin D_3, bone-breaking load, egg production, plasma P, 1233†
egg shell quality, vitamin D, 166*
adult selection, genetic merit, juvenile selection, selection response, 1655
barley, egg weight, feed utilization, metabolizable energy, 1306

behavior, cage shape, feed trough partitions, 1348†
broiler chick performance, sesbania seed, 11*
cage area, corticosterone, dominance rank, egg production, 399
cage shape, nipple waterer, waterer location, waterers per cage, 89*
cage type, feeder type, hen density, 102*
Cassia obtusifolia seed, coffee weed, nutrient fortification, 15*
chlorella, egg production, 84*
cholesterol, herring meal, omega-3 fatty acids, 99*
cystine, methionine, transsulfuration enzyme activity, 78*
D-Trp-6-LHRH, molt-induced, 19*
days without eggs, fasting periods, feather loss, peaks of production, 67*
dietary acidification, dietary alkalinization, renal function, urolithiasis, 1694
dietary protein, egg production, pullet dietary treatment, 62*
early protein restriction, egg type pullet, growth restriction, 54*
egg production, egg quality, field beans, vicine, 455
egg weight, feed efficiency, Fermacto®, methionine level, 842†
energy level, pullet diet, 63*
environmental stress, relative humidity, temperature, vitamin C, 67*
feed conversion, phase feeding, protein dietary regimens, 1447
force resting, iron, zinc, 3*
heat stress, hypothalamus, luteinizing hormone, luteinizing hormone releasing hormone, 79*
meal feeding, 125*
phosphorus, sodium aluminosilicate, 31*

Layer performance year round, Japan, open-sided houses, photoperiod, seasonal change, 1130
Layer replacement, egg production rate, multiple hen cages, 848†
Layer requirements, lysine, sulfur amino acids, 130*
Laying photoperiod, ahemeral photoperiod, egg production, rearing photoperiod, shell quality, 391
Lead, body weight, calcium, egg production, quail-Japanese, 81*
Lecithin, fat, phospholipids, shell gland, shell strength, 8*
Lecithovitellin reaction, egg yolk, food-poisoning bacteria, salted yolk paste, *Staphylococcus aureus*, 261
Leg abnormalities, body weight, feed conversion, niacin requirement, turkey poult, 760
broiler chick, isoleucine, leucine, valine, 85*
glucagon, insulin, lame *vs.* normal, turkey, 97*
Leg bones, body weight, leg muscles, shank width, turkey, 1660
Leg muscles, body weight, leg bones, shank width, turkey, 1660
Leghorn, amino acid(s), bioavailability, energy, guinea fowl, 145*
Leghorn cockerels, air velocity, cage environment, energy consumption, feed intake, 62*
Leghorn male, feeding time, meal feeding, plasma, uric acid, 345†
Lesions, hypothalamus, obesity, sexual dysfunction, 157*
Leucine, broiler chick, isoleucine, leg abnormalities, valine, 85*
Leukocyte distribution, antibody production, growth hormone, lymphoid organs, T_3,T_4 doses, 92*
Leupeptin, chick embryo, proteolysis, skeletal muscle cell culture, 87*
Ligand, cytochrome c, heme-complex, myoglobin, poultry meat color, 44*
Light environment, endocrine glands, light intensity, population density, turkey performance, 1518
Light intensity, abdominal fat, broiler, feed conversion, growth rate, 73*
broiler growth, 1239
behavior, broiler performance, 1020
endocrine glands, light environment, population density, turkey performance, 1518
locomotion, photostimulation, quail-Japanese, 120*
Light regimen, antibody production, immune response, thymidine uptake, white blood cell, 114*
turkey breeder performance, 173*
Light source, broiler diet, energy use, feed efficiency, weight gain, 47*
broiler growth, efficiency of production, 151*
broiler growth performance, fluorescent, incandescent, sodium vapor, 43
Lighting environment, feed form, floor space, turkey management, 127*
Lighting program, ahemeral lighting, egg production, egg weight, pullet, 176*
androgen, broiler, organ weight, restricted feeding programs, 145*
blackout rearing, body weight, broiler breeder performance, 2*
body weight, broiler breeder performance, egg production, 48*
broiler breeder, fat in diet, fertility, hatchability, 5*
broiler performance, feed restriction, skeletal diseases, 68*
diurnal rhythm, feather condition, layer, operant heat demand, 108*
energy level, layer diet, prolapse, pullet diet, 108*
pullet, self-select diets, sexual maturity, temperature, 152*
Limestone, calcium source, layer digestive system, particle size, solubility, 142*
particle size, solubility, 29*
Limestone solubility, calcium source, egg production, particle size, shell quality, 66*
Linkage relationship, barring, dermal melanin, recessive white skin, sex chromosome, 530
Linoleate, biotin deficiency, bone modeling, broiler chick, valgus-varus, 590
Linoleic acid, corn oil, egg mass, 118*
Lipase, abdominal fat, genetic correlation, lipoprotein, protein, 183
Lipase activity, 17β-estradiol dipropionate, pancreas, turkey poult, 84*
pancreas, turkey poult, yolk membrane, 1089
Lipid content, chicken patties, frying conditions, moisture content, shortening quality, 9*
Lipid fraction, carcass composition, energy estimation, phospholipids, protein content, 413
Lipid oxidation, antioxidant, mechanically deboned meat, oxidative stability, 270
fat sources, fatty acid profile, sausage, 91*
functional properties of myofibrils, myofibrils, storage time, water activity, 128*
Lipid stability, broiler meat quality, carcass weepage, vitamin E, 38*
Lipogenesis, ascorbic acid, broiler male chick, corticosterone implant, 1356†
broiler chick, compensatory growth, energy repletion, enzyme activities, 146*
lysine, protein, 146*

Lipoprotein, abdominal fat, genetic correlation, lipase, protein, 183
Listeria monocytogenes, broiler, colonization, 53*
cell wall antigen, enzyme immunoassay, 157*
cooked chicken, ELISA, 157*
Litter, rickets, turkey poult, 106*
Litter condition, body weight, broiler diet, triticale, 1743
floor type, turkey performance, 66*
Litter feeding, broiler litter, computer simulation, economic value, 1352†
Litter moisture, anticoccidial drugs, broiler, coccidial lesion scores, growth performance, 126*
niacinamide, stocking density, turkey poult performance, 24*
Litter treatment, ammonia, broiler production, 151*
Livability, caged layer, costs, culling procedures, management tool, 148*
Liver, bile acids, broiler chick, intestine, SEHCAT, 17*
dietary fat, hepatic tocopherol concentration, turkey, vitamin E, 1814†
Liver lipids, biotin, chick, fatty acids, *trans*/18:1, 45*
Local barley, broiler performance, corn substitution, Iraq rations, 122*
Location, broiler, intestine, Meckel's diverticulum, yolk stalk, 677†
Locomotion, light intensity, photostimulation, quail-Japanese, 120*
Luprosil, broiler performance, coliforms, *Escherichia coli*, 13*
Lutein, aflatoxin, carotenoids, pale bird syndrome, pharmacokinetics, 619, 1080
carotenoid composition, egg, serum, yolk, 608
Luteinizing hormone, follicle-stimulating hormone, granulosa cell, inhibin activity, 1625
glycoprotein hormone, hybridoma, monoclonal antibodies, turkey, 117*
heat stress, hypothalamus, layer performance, luteinizing hormone-releasing hormone, 79*
layer, ovarian follicle, steroidogenesis, 145*
Luteinizing hormone-releasing hormone, monoclonal antibody, quail, 110*
cyclic adenosine monophosphate, diacylglycerol, luteinizing hormone secretion, 101*
heat stress, hypothalamus, layer performance, luteinizing hormone, 79*
Luteinizing hormone secretion, cyclic adenosine monophosphate, diacylglycerol, luteinizing hormone-releasing hormone, 101*
Lymphocyte antigens, cytotoxic T cells, monoclonal antibodies, natural killer cells, 111*
Lymphocyte response, *B*-congenic chickens, flow cytometric analysis, Ia antigens, monoclonal antibody, 1525
Lymphoid organs, antibody production, growth hormone, leukocyte distribution, T_3,T_4 doses, 92*
Lymphokines, coccidiosis, *Eimeria tenella*, 105*
Lysine, abdominal fat, broiler, carcass measurement, feather meal, 111*
amino acid(s), broiler performance, methionine, protein requirements, 36*
growth rate, methionine, turkey tom performance, 86*
body weight, threonine, turkey poult, valine, 140*
broiler breast meat, broiler performance, methionine, 98*
broiler carcass quality, meat yield, 123*
broiler chick, lipogenesis, protein, 146*
diaminopimelic acid, microorganisms, rooster-intact, cecectomized or cannulated, 1345†
layer requirements, sulfur amino acids, 130*
Lysine assay, egg mass, egg production, layer, soybean meal, 49*
Lysine deficiency, amino acid deficiency, arginine deficiency, chick growth, chloride excess, 172*
chick growth, *Eimeria acervulina*, feed efficiency, 232
Lysine requirement, amino acid(s), broiler, carcass composition, 161*
protein source, two-slope broken line method, 53*
Lysine utilization, arginine utilization, chick, growth assay, monensin, 341†
Macrophages, Congo Red negative, Congo Red positive, *Escherichia coli*, 121*
selenium deficiency, vitamin E deficiency, 141*
Maduramicin, anticoccidial drugs, narasin, nicarbazin, salinomycin, 35*
broiler performance, halofuginone, monensin, nicarbazin, 32*
Magnesium, bone ash, broiler chick, ionophores, 59*
Magnesium deficiency, broiler chick, parathyroid, plasma minerals, 6*
Magnesium excess, bone, egg production, egg shell quality, feed intake, 97*
layer, phosphorus, plasma, 18*
Majors *vs.* nonmajors, poultry component in introductory course, student performance, 149*
Male chickens, antibody, *Brucella abortus*, heterophils, humoral immunity, 1775†
Male-feeding system, broiler breeder performance, egg production, fertility, hatchability, 48*
Malposition, day of incubation, egg turning, hatchability, high density plastic egg flat, 685
Mammalian embryo, chick embryo, shell-less culture, 55*
Management, broiler chick performance, semi-isolation brooding, weight gain, 96*
care of poultry, feed intake principles, 90*
Management tool, caged layer, costs, culling procedures, livability, 148*
Manganese, bioavailability, chick, 41*
Manganese methionine, bioavailability, chick, 18*
Manganese utilization, calcium excess, chick, phosphorus excess, 172*
Mannose sensitivity, chick, intestine, receptors, *Salmonella typhimurium*, 133*
Manure, hatch time, house fly, larvicide, thiourea, 407
Manure moisture, citrinin, diuresis, kidney function, ochratoxin A, 1034
Manure-cage layer, carbon sources, coliforms, fermentation, *Salmonella*, 74*
Marek's disease, embryo, hatching egg injection machine, incubation, vaccination, 110*
Marek's disease vaccine, public health, 141*
Marinade composition, exudate, salt soluble protein, tumbling, turkey meat, 162*
Marinated broiler, carcass size, moisture absorption, moisture retention, sodium pyrophosphate, 510†
Marinated chicken fillet, cooking method, moisture content, volatile analysis, yield, 128*
Marination, breast muscle, moisture binding, rigor, shear force, 175*
Market survey, antibiotic resistance, broiler carcass, *Salmonella*, 55*
Mash, egg type pullet, pellets *vs.* mash, temperature, weight gain, 375
Maxiban®, broiler male performance, heat stress, mortality, nicarbazin, 17*
Mazidagi raw rock phosphate, broiler diet, layer diet, phosphorus source, 154*

Meal feeding, body fat, chick, dietary energy, feed intake, 140*
feeding time, Leghorn male, plasma, uric acid, 345†
layer performance, 125*
Meat and bone meal, abdominal fat, broiler performance, metabolizable energy values, 294
apparent metabolizable energy, metabolizable energy, true metabolizable energy, 280
Meat characteristics, carcass yield, guinea fowl, 82*
Meat composition, cooking, meat source, turkey roast, 1810†
Meat flavor, beef sausage, fat content, quality attributes, turkey sausage, 1075
dehulling, turkey growth, white lupine, 596
Meat meal, amino acid(s), bioavailability, protein quality, 65*
Meat quality, broiler breast meat, electrical stunning, hot boning, 674†
Meat source, cooking, meat composition, turkey roast, 1810†
Meat yield, broiler carcass quality, lysine, 123*
turkey tom, 43*
Mechanically deboned meat, antioxidant, lipid oxidation, oxidative stability, 270
composition, functional properties, washing, 87*
kamaboko-like products, poultry meat color, yield, 112*
cooked gels, spray-dried egg white, water-washed, 73*
gel-strength, poultry meat color, surimi-wash process, 91*
Meckel's diverticulum, broiler, intestine, location, yolk stalk, 677†
Melatonin level, pineal gland, *rc* chickens, retina, serum, 66*
Membrane layers, egg, glycoprotein, ovomucin, vitelline membrane, 476
Mesotocin, arginine vasotocin, broiler, heat stress, surgery stress, 171*
Metabolic clearance rate, age, growth hormone, protein restriction, secretion rate, 120
Metabolic rate, conductive heat loss, heat stress, panting, roost temperature, 124*
Metabolism, Alimet®, broiler chick, DL-methionine, excreta, 12*
canthaxanthin, chick, hydroxyechinenone, isozeaxanthin, 787
Metabolites, blood chemistry, chick, organ weight, sterigmatocystin, 502
Metabolizable energy, abdominal fat, broiler performance, poultry by-product meal, 119*
acid-insoluble ash, amaranth, broiler, 641
animal by-product meal, protein, 72*
apparent metabolizable energy, meat and bone meal, true metabolizable energy, 280
barley, egg weight, feed utilization, layer performance, 1306
broiler performance, fat, grain sorghum, tannin content, 80*
Metabolizable energy assays, fasted rooster, feather meal, variances, 1188
Metabolizable energy intake, layer, prediction equation, 135*
Metabolizable energy utilization, carcass composition, energy, layer, temperature, 136*
Metabolizable energy values, abdominal fat, broiler performance, meat and bone meal, 294
Metachromatic staining, muscle fiber type, 982†
Metaphase chromosome, *B*-actin gene, gene mapping, *in situ* hybridization, 154*
Methionine, Alimet®, brown egg layer, egg mass, feed efficiency, 16*
excreta, 76*
amino acid(s), broiler performance, lysine, protein requirements, 36*
growth rate, lysine, turkey tom performance, 86*
bioavailability, digestibility, methionine hydroxy analog, 94*
broiler, methionine hydroxy analog, oxidation, 81*
broiler breast meat, broiler performance, lysine, 98*
broiler chick, digestibility, DL-methionine hydroxy analog, 14*
broiler performance, monensin, nicarbazin, salinomycin, 33*
choline, turkey poult, 9*
cystine, layer performance, transsulfuration enzyme activity, 78*
Methionine activity, corn-soy-gelatin basal diet, methionine assay, soybean meal, 88
Methionine assay, corn-soy-gelatin basal diet, methionine activity, soybean meal, 88
Methionine hydroxy analog, bioavailability, digestibility, methionine, 94*
broiler, methionine, oxidation, 81*
broiler chick, intestinal uptake, rat, 56*
Methionine level, broiler performance, sodium sulfate, 159*
egg weight, feed efficiency, Fermacto®, layer performance, 842†
Methionine requirement, age, layer, sulfur amino acid requirement, 170*
Methionine supplementation, choline supplementation, egg yolk, high performance liquid chromatography, phospholipids, 1717
Methylation, 5-azacytidine, prolactin, turkey, 86*
MHC genes, aneuploids, bursal lymphocyte population, 74*
Microbial contamination, broiler, chemical additives, poultry-processing water, *Salmonella*, 10*
Microbiological contamination, broiler, feather release, picking feathers, spray scalding, 12*
Microbiological growth support, futher processing, nutrient availability, particle size, turkey meat product form, 1269
Microorganisms, diaminopimelic acid, lysine, rooster-intact, cecectomized or cannulated, 1345†
Microwave cooking, broiler meat, *Salmonella*, 71*
Microwave radiation, broiler egg, immune response, incubation, 57*
chick incubation, reproductive performance, 57*
chlordecone, corticosterone, embryo, quail-Japanese, 60*
Migration, blood leukocyte, dopamine, norepinephrine, serotonin, 116*
Milo, bacteria, broiler carcass speck count, corn, intestinal spillage, 88*
Mineral anions, acid-base balance, broiler chick, tibial dyschondroplasia, 148*
Mineral level, embryonic mortality, fertility, 127*
Mineral level in eggs, fertility, mineral level in hen blood, 163*
Mineral level in hen blood, fertility, mineral level in eggs, 163*
Minimum time process, accelerated rigor, electric stimulation, nonaged tenderness, tenderness, 40*
Mitogen response, erythrocyte alloantigens, hemagglutination reactions, immunogenetic analysis, immuno-

globulin G, 1210
Mitogenesis, carrageenan, *in vitro*, peripheral blood lymphocytes, phytohemagglutinin-M, 33*
Mitogenphytohemagglutinin, broiler chick, cortisol, cutaneous basophil hypersensitivity, 669
Moisture, antifungal activity, gentian violet, heat, particle size, 58
Moisture absorption, carcass size, marinated broiler, moisture retention, sodium pyrophosphate, 510†
Moisture binding, breast muscle, marination, rigor, shear force, 175*
Moisture content, chicken patties, frying conditions, lipid content, shortening quality, 9*
cooking method, marinated chicken fillet, volatile analysis, yield, 128*
feed and water withdrawal, gastrointestinal tract, shear strength, water withdrawal, 845†
Moisture pick-up, broiler meat, cooking yields, heat stress, 27*
Moisture retention, carcass size, marinated broiler, moisture absorption, sodium pyrophosphate, 510†
Molt, body weight, feed restriction, temperature, 137*
D-Trp-6-LHRH, gonadotroph, layer, reproductive performance, 38*
egg production, photorefractoriness, thyroxine, turkey, 21*
egg shell quality, pimpling, 205
postmolt performance, temperature, 126*
Molt-induced, aluminum, egg production, quail-Japanese, zinc, 1157
D-Trp-6-LHRH, layer performance, 19*
duodenum, feed consumption, layer, zinc, 58*
egg shell strength, phospholipids, shell gland, 427
egg weight, layer, zinc, 5*
Monensin, abdominal fat pad, adipocyte size, broiler, withdrawal period, 339†
animal fat, body weight, broiler chick performance, feed intake, 452
anticoccidial drugs, chick, protein level, protein source, 652
anticoccidial efficacy, field trials, narasin, 1058
arginine utilization, chick, growth assay, lysine utilization, 341†
body weight control, broiler breeder, phenypropanolamine, 133*
broiler performance, halofuginone, maduramicin, nicarbazin, 32*
methionine, nicarbazin, salinomycin, 33*
chick weight gain, coccidiosis, duodenal pH, zeolite-A, 171*
Monoclonal antibodies, broiler, ELISA, *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, 116*
B-congenic chickens, flow cytometric analysis, Ia antigens, lymphocyte response, 1525
cytotoxic T cells, lymphocyte antigens, natural killer cells, 111*
glycoprotein hormone, hybridoma, luteinizing hormone, turkey, 117*
luteinizing hormone-releasing hormone, quail, 110*
Morphogenesis, embryogenesis, pigeon, secondary plate development, 865
Mortality, age, beak trimming, feed consumption, guinea fowl, 141†
broiler, feed restriction, heat stress, 2*
air sac lesions, broiler, feed efficiency, fructooligosaccharides, 46*
biogenic amines, duodenum, hypothalamus, turkey poult, 13*
broiler, carbonated water, feed intake, heat stress, 30*
feed restriction, heat stress, 226
broiler breeder performance, egg production, floor type, hatchability, 1414
rabbit excreta, 39*
broiler chick growth, gizzard erosion, gizzerosine, 1580
broiler male performance, heat stress, Maxiban®, nicarbazin, 17*
carcass preservation, dead bird disposal, preservation acid, 113*
chemical composition, culls, layer, 94*
Mucin layer, ceca, *Salmonella* attachment, 26*
Multiphasic analysis, growth curve, weight gain, 33
Multiple hen cages, egg production rate, layer replacement, 848†
Multiple stressors, behavior, caging system-experimental, chick performance, health, 1787†
Muscalure, black light devices, caged layer, electrocuting flies, flies, 871
Muscle characteristics, broiler breast muscle, broiler drumsticks, broiler growth, exercise, 1557
Muscle development, growth, turkey genotypes, 22*
Muscle fiber diameter, breast muscle, muscle fiber type, 908
Muscle fiber type, breast muscle, muscle fiber diameter, 908
histochemistry, skeletal muscle, 973
metachromatic staining, 982†
Muscle growth, breast muscle, broiler, exercise patterns, thigh, 1708
gene interactions, genetics, muscular dystrophy, 367
myogenic, satellite cell, turkey, 121*
Muscle pH, delayed glycolysis, electric stunning, protein solubility, turkey muscle, 1062
Muscle tenderness, broiler, pH, post-mortem temperature, sarcomere length, 54*
Muscular dystrophy, gene interactions, genetics, muscle growth, 367
Mutation, Fawn, genetics, plumage color, quail-Japanese, 127*
Myco Curb®, broiler carcass, chemical preservative, feed, *Salmonella*, 1225†
Mycoplasma gallisepticum, antibody, *Mycoplasma synoviae*, 58*
broiler, ELISA, monoclonal antibodies, *Mycoplasma synoviae*, 116*
cloning, *Escherichia coli*, 26*
egg production, egg shell quality, layer, 5*
heat stress, layer, 34*
Mycoplasma species, SDS-PAGE analysis, 116*
***Mycoplasma gallisepticum* identification**, flow cytometer, *Mycoplasma synoviae* identification, 25*
***Mycoplasma* species**, *Mycoplasma gallisepticum*, SDS-PAGE analysis, 116*
Mycoplasma synoviae, antibody, *Mycoplasma gallisepticum*, 58*
broiler, ELISA, monoclonal antibodies, *Mycoplasma gallisepticum*, 116*
***Mycoplasma synoviae* identification**, flow cytometer, *Mycoplasma gallisepticum* identification, 25*
Mycotoxin, aflatoxin, aluminas, aluminosilicate, detoxification, 243
broiler male chick, synergy, T-2 toxin, 1418
age, blood chemistry, broiler, ochratoxin A, 1139
copper, dyschondroplasia, fusarochromanone, immunity, 174*
Myoelectrical activity, oviposition, uterus, vagina, 1615
Myofibrils, functional properties of myofibrils, lipid oxida-

tion, storage time, water activity, 128*
Myogenic, muscle growth, satellite cell, turkey, 121*
Myoglobin, cytochrome c, heme-complex, ligand, poultry meat color, 44*
heme-complex, hemoglobin, pH, poultry meat color, 44*
Myopathies, growth, turkey muscle, 173*
Myosin, actin, avian skeletal muscle, collagen, N^{τ}-methylhistidine, 1593
N^{τ}-methylhistidine, actin, avian skeletal muscle, collagen, myosin, 1593
Naked neck, body weight, heat stress, 13*
fertility, reproductive traits, temperature, 176*
Naked neck *vs.* normal, body temperature, heat stress, 818
Naked oats, chemical composition, layer diet, weathering, 115*
Naked oats pricing, broiler, economics, layer, turkey, 10*
Narasin, anticoccidial drugs, maduramicin, nicarbazin, salinomycin, 35*
anticoccidial efficacy, battery cage trials, *Eimeria*, ionophorous antibiotic, 1043
Eimeria, floor pen trials, ionophorous antibiotic, 1050
field trials, monensin, 1058
bacitracin, body weight, broiler, roxarsone, 103
broiler performance, coccidiosis, heat stress, nicarbazin, 248
clearance kinetics, layer, 8*
Natural killer cells, B-cells, growth hormone, T-cells, T_3,T_4 doses, 93*
cytotoxic T cells, lymphocyte antigens, monoclonal antibodies, 111*
Nematodes, anthelmintics, efficacy, 41*
Nervousness, egg production, feathering, pullet, rearing environment, 1400
Neural cell adhesion molecule, antibody perturbations, chick, regenerating nerve fibers, 143*
Neurological behavior, genetic lines, immunity, intracerebral inoculation, Rous sarcoma virus, 1123
Neuropeptide, chick, feed intake, hypothalamus, 107*
Newcastle disease virus, antibody response, egg production, *Pasteurella multocida*, turkey, 1372
Niacin, tryptophan, trytophan:niacin conversion, turkey poult, 148*
Niacin requirement, body weight, feed conversion, leg abnormalities, turkey poult, 760
Niacin supplementation, corn-soybean meal diet, fertility, hatchability, turkey breeder performance, 336†
Niacinamide, litter moisture, stocking density, turkey poult performance, 24*
Nicarbazin, acid-base balance, broiler, heat stress, 3*
anticoccidial drugs, coccidiosis, halofuginone, polyether ionophores, 31*
maduramicin, narasin, salinomycin, 35*
arprinocid, broiler resistance, *Eimeria tenella*, glycamide, 164*
broiler male performance, heat stress, Maxiban®, mortality, 17*
broiler performance, coccidiosis, heat stress, narasin, 248
halofuginone, maduramicin, monensin, 32*
methionine, monensin, salinomycin, 33*
Nicotinamide, heme-complex, poultry meat color, pyridine, turkey breast meat, 44*
Nipple waterer, broiler chick, flow rate, 137*
cage shape, layer performance, waterer location, waterers per cage, 89*
Nitrogen retention, broiler breeder male, diet, standard *vs.* all-corn diet, 144*
Nonaged tenderness, accelerated rigor, electric stimulation, minimum time process, tenderness, 40*
Nonestrified fatty acids, heparin, turkey, very low density lipoprotein, 51*
Nonlinear models, egg production cycles, egg production models, 213
Nonspecific nitrogen source, abdominal fat deposition, broiler, feather meal, finishing period, 300
Norepinephrine, blood leukocyte, dopamine, migration, serotonin, 116*
catecholamines, dopamine, epinephrine, 75*
Northern fowl mite, antibody, immune response, layer, Western blot analysis, 549
bovine serum albumin injection, host immunity, rocket electrophoresis, two-dimensional immunoelectrophoresis, 1015
Nucleotide sequence, cloning, growth hormone, 107*
Nuisance abatement, grower organization, industry regulation, peer pressure, 169*
Numida melagris, genetics, guinea fowl, plumage color, 158*
Nutrient availability, futher processing, microbiological growth support, particle size, turkey meat product form, 1269
Nutrient fortification, *Cassia obtusifolia* seed, coffee weed, layer performance, 15*
Nutrient restriction, abdominal fat, broiler, 6*
Nutrition, breed effect, roaster weight, temperature regimen, 96*
progeny performance, turkey breeder, 170*
Nutritional requirements, growth, quail-Japanese, selection, 22*
Nutritional status, egg assay for riboflavin, layer, riboflavin, 160*
Nutritive value, amaranth, apparent metabolizable energy, broiler chick growth, 1166
Oat hulls, aerobic fermentation, broiler chick, poultry droppings, 52*
Obesity, hypothalamus, lesions, sexual dysfunction, 157*
Ochratoxin A, age, blood chemistry, broiler, mycotoxin, 1139
body weight, broiler chick performance, organ weight, phenylalanine, 90*
broiler male chick, deoxynivalenol, organ weight, 253
citrinin, diuresis, kidney function, manure moisture, 1034
pullet, renal function, 90*
egg, quail-Japanese, tissue distribution, whole body autoradiography, 707
Odor, dimethyl sulfide, egg products, spoilage, uracil, 123*
Off-flavor, chicken patties, reheating methods, sensory profiling, thiobarbituric acid, 112*
Oil, confounding experimental results, energy supplementation, growth assay, sand, 169*
Oligosaccharides, digestibility, soybean meal, transit time, 109*
Omega-3 fatty acids, cholesterol, egg, full-fat canola seed, serum, 129*
herring meal, layer performance, 99*
Open-sided houses, Japan, layer performance year round, photoperiod, seasonal change, 1130
Operant heat demand, diurnal rhythm, feather condition, layer, lighting program, 108*
Optimization, broiler growout, computer program, simulation program, temperature, 534
Organ culture, chick, insulin, pancreas, 101*

Organ proportions, carcass quality, compensatory growth, protein undernutrition, turkey, 1760
Organ weight, androgen, broiler, lighting program, restricted feeding programs, 145*
blood chemistry, chick, metabolites, sterigmatocystin, 502
body weight, broiler chick performance, ochratoxin A, phenylalanine, 90*
broiler male chick, deoxynivalenol, ochratoxin A, 253
chick growth, deoxynivalenol, wheat contamination, 1778†
Organic acids, broiler diet, calcium formate, formic acid, fumaric acid, 1178
Ovarian cells, prolactin, steroidogenesis, turkey, 95*
Ovarian follicle, 17α-hydroxylase, hypophysectomy, quail-Japanese, theca cell, 1636
layer, luteinizing hormone, steroidogenesis, 145*
Ovarian follicle growth, guinea fowl, pigeon, 1783†
Ovarian growth, estradiol, pregnant mare serum gonadotropin, prolactin, turkey, 319
Ovarian tissue, antibiotics, bovine luteinizing hormone, steroidogenesis, 1486†
Oven type, cooking time, turkey roast size, 139*
Oviposition, double oviduct, experimental surgery, fertility, 1465
myoelectrical activity, uterus, vagina, 1615
Oviposition rate, defective eggs, genetic associations, shell quality, shell-less egg, 123*
Ovomucin, egg, glycoprotein, membrane layers, vitelline membrane, 476
Oxidation, broiler, methionine, methionine hydroxy analog, 81*
Oxidative stability, antioxidant, lipid oxidation, mechanically deboned meat, 270
bacterial flora, broiler meat, storage time, 2*
Oxygen affinity, 2,3-bisphosphoglycerate, erythrocytes, inositol pentakisphosphate, phosphate, 100*
Oxygen tension, embryonic mortality, high altitude, incubation temperature, turkey egg hatchability, 956
Ozonated water, formaldehyde, hatching egg sanitizer, hydrogen peroxide, quaternary ammonium compound, 155*
hatching egg sanitizer, hydrogen peroxide, quaternary ammonium compound, 57*
P150 merozoite protein, antibody response, *B*-congenic chickens, *Eimeria acervulina*, recombinant coccidial antigens, 111*
Package size, consumer preference, consumer preference, poultry products, 50*
Packaged carcasses, broiler, chilling time, glycol solution, temperature, 154*
Pad cooling, broiler, evaporative cooling, fan cooling, ventilation, 98*
Pale bird syndrome, aflatoxin, carotenoids, lutein, pharmacokinetics, 619, 1080
Palisade layer, abnormal egg shell structure, calcium reserve, egg shell structure, 77*
Pancreas, 17β-estradiol dipropionate, lipase activity, turkey poult, 84*
chick, insulin, organ culture, 101*
lipase activity, turkey poult, yolk membrane, 1089
Pancreas weight, body weight, soybean meal, turkey, urease index, 15*
Pancreatic lipase, glycerol trioleate, turkey, 85*
Pancreatic polypeptide, biliary secretion, layer, pancreatic secretion, 126
Pancreatic secretion, biliary secretion, layer, pancreatic polypeptide, 126
Panting, conductive heat loss, heat stress, metabolic rate, roost temperature, 124*
Parathyroid, broiler chick, magnesium deficiency, plasma minerals, 6*
Particle size, antifungal activity, gentian violet, heat, moisture, 58
calcium source, egg production, limestone solubility, shell quality, 66*
layer digestive system, limestone, solubility, 142*
limestone, solubility, 29*
futher processing, microbiological growth support, nutrient availability, turkey meat product form, 1269
Parts composition, body weight, carcass yield, turkey, 153*
Passive immunity, antibiotics, antiurease antibody, chick growth, jackbean urease enzyme, 434
Pasteurella multocida, antibody response, egg production, Newcastle disease virus, turkey, 1372
turkey, 149*
attenuated vaccine, *Escherichia coli*, 113*
dietary protein restriction, fowl cholera, humoral immunity, 135*
disease challenge, egg production, turkey, 1807†
fowl cholera, KSCN extract, turkey, vaccination, 26*
Pathogenesis, infectious bronchitis virus, tissue tropism, 60*
Pavo cristatus, genetics, peafowl, plumage color mutation, sex-linked gene, 158*
Peafowl, genetics, *Pavo cristatus*, plumage color mutation, sex-linked gene, 158*
Peaks of production, days without eggs, fasting periods, feather loss, layer performance, 67*
Peanut skins, chick diet, copper, weight gain, zinc, 51*
Peck order, caged layer, despotism, dominance, welfare, 1008
***Pectoralis* muscle**, chewiness, cooking method, deboning time, hardness, 112*
denervation, 60*
Peer pressure, grower organization, industry regulation, nuisance abatement, 169*
Pellet binder, blood chemistry, formaldehyde, layer, urea, 37*
Pelleting, Aureomycin, egg production, pullet growth, sunflower seeds, 445
broiler, feed efficiency, sunflower meal, 175*
canola seed, full-fat canola seed, seed processing methods, turkey, 1731
Pellets, barley, corn texture, grit, turkey performance, 101*
energy, protein level sequence, turkey tom performance, 170*
Pellets *vs.* mash, beak trimming, body weight, broiler roaster growth, feed intake, 1514
egg type pullet, mash, temperature, weight gain, 375
Peptide histidine isoleucine, prolactin-releasing factor, turkey, 141*
Peptides, catecholamines, chicken, feed intake, turkey, 74*
Performance index, chloride level, layer, phosphorus level, potassium level, 131*
Peripheral blood lymphocytes, carrageenan, *in vitro*, mitogenesis, phytohemagglutinin-M, 33*
Perivitelline layer, acrosome reaction, cock spermatozoa, fertilization, *in vitro* incubation, 1770
Perlite, abdominal fat pad, broiler performance, wheat, 33*
Peroxide, broiler performance, ethoxyquin, feed preservative, rancidity, 1725
pH, adenosine triphosphate, electric stimulation, poultry carcass, tenderness, 106*
broiler, muscle tenderness, post-mortem temperature,

sarcomere length, 54*
broiler breast meat, denervation, lactate, R value, 117*
heme-complex, hemoglobin, myoglobin, poultry meat color, 44*

Phagocytosis, antibody response, histocompatibility complex, immunocompetence, T cell-mediated response, 989

Pharmacokinetics, aflatoxin, carotenoids, lutein, pale bird syndrome, 619, 1080

Phase feeding, feed conversion, layer performance, protein dietary regimens, 1447

Pheasant breeder, brewers dried grains, egg production, fertility, hatchability, 138*

Phenylalanine, body weight, broiler chick performance, ochratoxin A, organ weight, 90*

Phenypropanolamine, body weight control, broiler breeder, monensin, 133*

Phosphate, 2,3-bisphosphoglycerate, erythrocytes, inositol pentakisphosphate, oxygen affinity, 100*

Phospholipids, biotin, broiler chick, fatty acids, 171*
carcass composition, energy estimation, lipid fraction, protein content, 413
choline supplementation, egg yolk, high performance liquid chromatography, methionine supplementation, 1717
egg shell strength, molt-induced, shell gland, 427
fat, lecithin, shell gland, shell strength, 8*

Phosphorus, bioavailability, body weight, toe ash, turkey, 96
blood levels, calcium, layer, shell quality, 68*
bone, layer, magnesium excess, plasma, 18*
broiler chick diet, by-product sources, precipitated bone phosphate, 1302
calcium, chloride, tibial dyschondroplasia, zeolite, 1436
layer performance, sodium aluminosilicate, 31*

Phosphorus absorption, calcium, layer, phosphorus excretion, 440

Phosphorus availability, bone ash, broiler, phosphorus sources, 152*
bone meal source, broiler performance, broiler strain, environmental temperature, 27*

Phosphorus excess, calcium excess, chick, manganese utilization, 172*

Phosphorus excretion, calcium, layer, phosphorus absorption, 440

Phosphorus level, chloride level, layer, performance index, potassium level, 131*
egg production, Ethacal® feed component, layer, 86*

Phosphorus metabolism, aluminum, broiler chick, vitamin D, 19*

Phosphorus requirement, calcium requirement, rickets, turkey poult, 150*

Phosphorus source, broiler diet, layer diet, Mazidagi raw rock phosphate, 154*

Phosphorus sources, bone ash, broiler, phosphorus availability, 152*

Photoperiod, color-filtered light, growth, turkey performance, 1505
feather color, quail-Japanese, testicular development, 1098†
Japan, layer performance year round, open-sided houses, seasonal change, 1130

Photorefractoriness, egg production, molt, thyroxine, turkey, 21*

Photostimulation, hypothalamus, prolactin, turkey, 138*
light intensity, locomotion, quail-Japanese, 120*

Physical manipulation, egg yolk, salt, shaking, texture, 1264

Phytohemagglutinin-M, carrageenan, *in vitro*, mitogenesis, peripheral blood lymphocytes, 33*

Picking feathers, broiler, feather release, microbiological contamination, spray scalding, 12*

Pigeon, embryogenesis, morphogenesis, secondary plate development, 865
guinea fowl, ovarian follicle growth, 1783†

Pigment source, broiler performance, broiler pigmentation, xanthophyll, 544

Pimpling, egg shell quality, molt, 205

Pineal gland, melatonin level, *rc* chickens, retina, serum, 66*

Plant feedstuffs, amino acid(s), bioavailability, *in vitro* digestion, true amino acid availability, 78

Plaque-forming cell transfer, bursal cell transfer, dopamine, immune response, sheep red blood cells, 88*

Plasma, body weight, genetic correlation, growth hormone, heritability, 1781†
bone, layer, magnesium excess, phosphorus, 18*
erythrocytes, hemolysis, t-butyl hydroperoxide, uric acid, 1632
feeding time, Leghorn male, meal feeding, uric acid, 345†

Plasma calcium, broiler breeder, feeding regimen, plasma phosphorus, 56*

Plasma corticosterone, body weight, broiler breeder pullet, feed restriction, plasma glucose, 809

Plasma free arachidonic acid, broiler, feed intake, 144*

Plasma free fatty acids, broiler, feed intake, force feeding, plasma ketones, 30*

Plasma glucose, body weight, broiler breeder pullet, feed restriction, plasma corticosterone, 809

Plasma hormone level, broiler, growth hormone, insulin-like growth factor-1, 117*

Plasma ketones, broiler, feed intake, force feeding, plasma free fatty acids, 30*

Plasma lipids, broiler, carcass fat, 129*

Plasma minerals, broiler chick, magnesium deficiency, parathyroid, 6*

Plasma P, 1,25-dihydroxyvitamin D_3, bone-breaking load, egg production, layer performance, 1233†

Plasma phosphorus, broiler breeder, feeding regimen, plasma calcium, 56*

Plasma thyroxine, dietary protein, quail-Japanese, thiouracil, 28*

Plasminogen activator, granulosa cell, growth factors, layer, 165*

Plumage color, black-tailed red pattern, dark brown gene, gene interactions, Villafranquina chickens, 351
Fawn, genetics, mutation, quail-Japanese, 127*
genetics, guinea fowl, *Numida melagris*, 158*

Plumage color mutation, genetics, *Pavo cristatus*, peafowl, sex-linked gene, 158*

Poly-L-glutamic acid, fertility, 82*

Polyether ionophores, anticoccidial drugs, coccidiosis, halofuginone, nicarbazin, 31*

Polypropylene container, broken-out eggs, refrigerated storage, 138*

Population density, endocrine glands, light environment, light intensity, turkey performance, 1518

Postinfection, anticoccidial efficacy, BAY Vi 9142, coccidiosis, toltrazuril, 122*

Postmolt performance, molt, temperature, 126*

Post-mortem aging, broiler breast muscle, cooking method, deboning time, texture profile, 23*

Post-mortem metabolism, boning time, broiler breast fillets, electric stimulation, tenderness, 150*

Post-mortem temperature, broiler, muscle tenderness, pH, sarcomere length, 54*

Potassium chloride, broiler performance, fasting, heat stress, weight gain, 36*
Potassium level, chloride level, layer, performance index, phosphorus level, 131*
Poultry by-product meal, abdominal fat, broiler performance, metabolizable energy, 119*
amino acid bioavailability, cecectomy, digestibility, protein quality, 97*
Poultry carcass, adenosine triphosphate, electric stimulation, pH, tenderness, 106*
Poultry component in introductory course, majors *vs.* nonmajors, student performance, 149*
Poultry droppings, aerobic fermentation, broiler chick, oat hulls, 52*
Poultry fat, broiler chick diet, fat blends, yellow grease, 27*
Poultry industry, extension poultry specialist role, extension program, poultry programs of the future, 883
Poultry inspection, Food Safety and Inspection Service, *Salmonella* contamination in poultry, *Salmonella* control, 944
Poultry majors, FFA and 4-H, poultry programs of the future, student recruitment, youth programs, 886
Poultry meat color, broiler breast meat, electric stimulation, hot boning, texture, 1536
composition, kamaboko-like products, mechanically deboned meat, yield, 112*
cytochrome c, heme-complex, ligand, myoglobin, 44*
gel-strength, mechanically deboned meat, surimi-wash process, 91*
heme-complex, hemoglobin, myoglobin, pH, 44*
nicotinamide, pyridine, turkey breast meat, 44*
Poultry-processing plant, flow volume measurement, water conservation, 119*
Poultry-processing water, bactericide, lactic acid, *Salmonella*, 69*
BOD removal, COD removal, hydrogen peroxide, wastewater treatment, 24*
broiler, chemical additives, microbial contamination, *Salmonella*, 10*
broiler carcass, detection, DNA probe, *Salmonella*, 80*
Poultry production, energy needs, feed intake, 115*
Poultry products, consumer preference, consumer preference, package size, 50*
irradiation, *Salmonella*, sensory quality, shelf life, 36*
Poultry programs of the future, business aspect of poultry industry, curriculum, scientific aspect of poultry industry, student recruitment, 879
extension poultry specialist role, extension program, poultry industry, 883
extension program, poultry students, teaching poultry, 878
FFA and 4-H, poultry majors, student recruitment, youth programs, 886
poultry research, poultry science departments, poultry students, 890
Poultry research, poultry programs of the future, poultry science departments, poultry students, 890
Poultry science, recruitment methods, undergraduate student, 160*
Poultry Science Association, Constitution and by-laws, 161
Poultry science departments, poultry programs of the future, poultry research, poultry students, 890
Poultry students, extension program, poultry programs of the future, teaching poultry, 878
poultry programs of the future, poultry research, poultry science departments, 890
Precipitated bone phosphate, broiler chick diet, by- product sources, phosphorus, 1302
Precocious puberty, chick, semen production, tamoxifen, 1642†
Prediction equation, broiler, water consumption, water:feed ratio, 131*
layer, metabolizable energy intake, 135*
Pregnant mare serum gonadotropin, estradiol, ovarian growth, prolactin, turkey, 319
Prelay pullet, bone, calcium level, egg shell weight, 98*
Preservation acid, carcass preservation, dead bird disposal, mortality, 113*
Probiotic, body weight, broiler, *Streptococcus faecium*, 132*
Processing plant, accident rate, 63*
broiler, chiller water, diatomaceous earth, filtration, 155*
Processing treatment, breast muscle, thigh muscle, turkey Canadian bacon, 162*
Processing water, (see poultry-processing water)
Processing yield, amaranth, body weight, broiler male performance, fat pad, 743
broiler, corn, fatty acids, sorghum, 130*
vitamin C, 85*
Productivity, broiler performance, growout process, quality, 4*
quality, 4*
Profitability, broiler grower, interregional competition, simulation model, 89*
Progeny performance, nutrition, turkey breeder, 170*
Progesterone, broody, egg production, egg quality, turkey, 72*
Prokaryotic expression vector, *Eimeria*, immune response, recombinant DNA, 100*
Prolactin, 5-azacytidine, DNA methylation, gene expression, turkey, 1794†
methylation, turkey, 86*
broody hen management, turkey, 128*
electric stimulation, serotonin, turkey, 93*
estradiol, ovarian growth, pregnant mare serum gonadotropin, turkey, 319
hypothalamus, photostimulation, turkey, 138*
ovarian cells, steroidogenesis, turkey, 95*
Prolactin-releasing factor, peptide histidine isoleucine, turkey, 141*
Prolapse, energy level, layer diet, lighting program, pullet diet, 108*
Propylthiouracil, egg production, pullet, sexual maturity, 107*
Prostaglandin, adrenocortical cells, corticosteroidogenesis, cyclic adenosine monophosphate production, 105*
Prostaglandin-E_2, corticosterone, quail-Japanese, stress, 32*
Protection antigen, antibody level, coccidiosis, immunization, 147*
Protein, abdominal fat, broiler, 7*
genetic correlation, lipase, lipoprotein, 183
amino acid(s), fat, turkey, 4*
animal by-product meal, metabolizable energy, 72*
animal fat, broiler performance, 80*
body composition, energy, turkey breeder performance, 17*
body weight, dietary fat, feed efficiency, turkey male, 1281
broiler chick, lipogenesis, lysine, 146*
dark turkey meat, sodium salts, storage, 103*
egg production, energy, pullet growth, 62*
energy, feed and water intake, feed efficiency, quail-Japanese growth, 1111

fingerprinting, infectious bronchitis virus, ribonucleic acid, 61*
soybean meal, toasting temperature, trypsin inhibitor, urease index, 20*
Protein and energy, growth rate, heritability, quail-Japanese, selection differentials, 519
Protein content, carcass composition, energy estimation, lipid fraction, phospholipids, 413
Protein degradation, age, breast muscle, broiler, protein synthesis, 135*
Protein dietary regimens, feed conversion, layer performance, phase feeding, 1447
Protein level, anticoccidial drugs, chick, monensin, protein source, 652
energy level, pullet, self-select diets, sexual maturity, 153*
fatty liver hemorrhagic syndrome, glutathione supplementation, quail-Japanese, 159*
Protein level sequence, energy, pellets, turkey tom performance, 170*
Protein malnutrition, adrenocortical cells, corticosterone production, 64*
Protein quality, amino acid bioavailability, cecectomy, digestibility, poultry by-product meal, 97*
amino acid(s), bioavailability, meat meal, 65*
Protein requirements, amino acid(s), broiler performance, lysine, methionine, 36*
Protein restriction, age, growth hormone, metabolic clearance rate, secretion rate, 120
body weight, feed efficiency, feed intake, turkey male performance, 130*
Protein solubility, broiler chick, soybean meal, urease activity, 11*
delayed glycolysis, electric stunning, muscle pH, turkey muscle, 1062
Protein source, anticoccidial drugs, chick, monensin, protein level, 652
lysine requirement, two-slope broken line method, 53*
Protein synthesis, age, breast muscle, broiler, protein degradation, 135*
Protein undernutrition, body composition, compensatory growth, turkey performance, 1750
carcass quality, compensatory growth, organ proportions, turkey, 1760
Proteolysis, chick embryo, leupeptin, skeletal muscle cell culture, 87*
Proximate analysis, ash, bomb calorimeter, feedstuffs, 513†
broiler breast muscle, glycogen content, hot boning, lactate content, 1702
Psittacine, *Clostridium*, ulcerative enteritis, 75*
Public health, Marek's disease vaccine, 141*
Pullet, acid-base balance, calcium level, urolithiasis, 90*
ahemeral lighting, egg production, egg weight, lighting program, 176*
beak trimming, behavior, feed consumption, weight gain, 109*
citrinin, ochratoxin A, renal function, 90*
egg production, feathering, nervousness, rearing environment, 1400
propylthiouracil, sexual maturity, 107*
energy level, protein level, self-select diets, sexual maturity, 153*
lighting program, self-select diets, sexual maturity, temperature, 152*
Pullet chick, behavior, body weight, rearing density, 47*
dietary calcium, infectious bronchitis virus, kidney function, urolithiasis, 1250
Pullet diet, energy level, layer diet, lighting program, prolapse, 108*
layer performance, 63*
Pullet dietary treatment, dietary protein, egg production, layer performance, 62*
Pullet growth, Aureomycin, egg production, pelleting, sunflower seeds, 445
egg production, energy, protein, 62*
Pullet strain, glomerular size, kidney anatomy, watering device, 31*
Pulsatile, broiler pullet, carcass composition, growth hormone, 168*
growth hormone, insulin, T_3,T_4, 168*
Pulverized and particle size limestone, acid-base balance, gypsum, layer diet, shell quality, 20*
Purification, biological activity, immune response, interleukin-1, 104*
Purification variables, breast muscle, cooking method, thiamin content, 72
Pyridine, heme-complex, nicotinamide, poultry meat color, turkey breast meat, 44*
Quaifenesin in diet, added fat in diet, reproductive performance, turkey hen, 114*
Quail, luteinizing hormone-releasing hormone, monoclonal antibody, 110*
Quail-Japanese, 17α-hydroxylase, hypophysectomy, ovarian follicle, theca cell, 1636
abdominal fat, beta agonist, eviscerated yield, feeding regimen, 120*
aluminum, egg production, molt-induced, zinc, 1157
atherosclerosis, cholesterol, exercise, 161*
genetic selection, 51*
body density, divergent selection, shank length, 88*
body weight, calcium, egg production, lead, 81*
busulfan, chemosterilant, 93*
chick, butylated hydroxyanisole, cysteine, lathyrism, toxicity, 153†
chlordecone, corticosterone, embryo, microwave radiation, 60*
control populations, egg production, egg weight, hatchability, 357
corticosterone, feed restriction, testosterone, 121*
heritability, stress, 25
prostaglandin-E_2, stress, 32*
dietary protein, plasma thyroxine, thiouracil, 28*
DNA modifications, egg yolk protein precursor, vitellogenin gene, 70*
dust-bathing activity, egg production, egg quality, fear-related response, 1363
egg, ochratoxin A, tissue distribution, whole body autoradiography, 707
egg weight loss, embryonic development, embryonic mortality, thiouracil, 28*
electron microscopy, epididymis, gossypol acetic acid, testis, 961
fatty liver hemorrhagic syndrome, glutathione supplementation, protein level, 159*
Fawn, genetics, mutation, plumage color, 127*
feather color, photoperiod, testicular development, 1098†
growth, nutritional requirements, selection, 22*
growth rate, heritability, protein and energy, selection differentials, 519
light intensity, locomotion, photostimulation, 120*
Quail-Japanese growth, energy, feed and water intake, feed efficiency, protein, 1111

Quality, broiler performance, growout process, productivity, 4*
productivity, 4*
Quality attributes, beef sausage, fat content, meat flavor, turkey sausage, 1075
Quantitative evaluation method, varus-angular bone deformity, 1647†
Quantitative traits, broiler breeder, dietary energy, selection, 6*
Quaternary ammonium compound, formaldehyde, hatching egg sanitizer, hydrogen peroxide, ozonated water, 155*
hatching egg sanitizer, hydrogen peroxide, ozonated water, 57*
R value, broiler breast meat, denervation, lactate, pH, 117*
broiler breast muscle, rigor development, sarcomere length, 635
R_2 alloantisera, avian leukosis virus, endogenous virus, hemagglutination, 50*
Rabbit excreta, broiler breeder performance, egg production, mortality, 39*
egg production, feed conversion, fertility and hatchability, semen production, 168*
Rabbits, cannibalism, chronic coccidiosis, congenital anomalies, ear, 102*
Radioreceptor assay, follicle-stimulating hormone, turkey, 95*
Rancidity, broiler performance, ethoxyquin, feed preservative, peroxide, 1725
Rapeseed meal, antinutritive effects, cecectomy, layer, 61*
Rat, broiler chick, intestinal uptake, methionine hydroxy analog, 56*
Rate of return, broiler, contract grower, economic analysis, 1804†
***Rc* chickens**, melatonin level, pineal gland, retina, serum, 66*
Rearing density, behavior, body weight, pullet chick, 47*
Rearing environment, egg production, feathering, nervousness, pullet, 1400
Rearing photoperiod, ahemeral photoperiod, egg production, laying photoperiod, shell quality, 391
Receptors, chick, intestine, mannose sensitivity, *Salmonella typhimurium*, 133*
Recessive white skin, barring, dermal melanin, linkage relationship, sex chromosome, 530
Recombinant coccidial antigens, antibody response, *B*-congenic chickens, *Eimeria acervulina*, p150 merozoite protein, 111*
coccidiosis, *Eimeria*, immune response, 68* ELISA, 103*
Recombinant DNA, chick, infectious bronchitis virus, viral diagnosis, 69*
coccidiosis, *Eimeria tenella*, vaccine, 72*
Eimeria, immune response, prokaryotic expression vector, 100*
Recruitment methods, poultry science, undergraduate student, 160*
Refrigerated storage, broken-out eggs, polypropylene container, 138*
Regenerating nerve fibers, antibody perturbations, chick, neural cell adhesion molecule, 143*
Regression equations, amino acid(s), crude protein, sorghum, 40*
Reheating methods, chicken patties, off-flavor, sensory profiling, thiobarbituric acid, 112*
sensory evaluation, storage time, thiobarbituric acid values, 736
Relatedness between individuals, DNA fingerprinting, 92*
Relative humidity, environmental stress, layer performance, temperature, vitamin C, 67*
Renal function, citrinin, ochratoxin A, pullet, 90*
dietary acidification, dietary alkalinization, layer performance, urolithiasis, 1694
Reproduction, body weight, dwarf Leghorns, sexual maturity, 143*
Reproductive performance, added fat in diet, quaifenesin in diet, turkey hen, 114*
age at sexual maturity, dietary protein, dwarf broiler breeder performance, hatchability, 219
chick incubation, microwave radiation, 57*
D-Trp-6-LHRH, gonadotroph, layer, molt, 38*
dietary protein, energy, triticale, turkey semen, 1619
semen production, thyroidectomy, turkey, 110*
Reproductive traits, fertility, naked nec^1, temperature, 176*
Resistance, genetic selection, growth, heat stress, 83*
Respiratory alkalosis, broiler, hyperthermia, hypocalcemia, lactate, 132*
Restricted feeding programs, androgen, broiler, lighting program, organ weight, 145*
Retina, melatonin level, pineal gland, *rc* chickens, serum, 66*
Retrovirus vector, chicken embryo, gene transfer, tissue-specific expression, 34*
chloramphenicol acetyltransferase, gene transfer, tissue-specific expression, 156*
gene transfer, selectable marker, 42*
Riboflavin, broiler chick, glutathione peroxidase, glutathione reductase, selenium, 149*
egg assay for riboflavin, layer, nutritional status, 160*
Riboflavin requirement, broiler chick, corn-soybean meal diet, 794
turkey poult, 32*
Ribonucleic acid, fingerprinting, infectious bronchitis virus, protein, 61*
Rickets, calcium requirement, phosphorus requirement, turkey poult, 150*
litter, turkey poult, 106*
Rigor, breast muscle, marination, moisture binding, shear force, 175*
Rigor development, aging temperature, broiler breast meat, 1147
broiler breast muscle, R value, sarcomere length, 635
Rigor mortis, broiler muscle, chilling, fiber type, 150*
Risk assessment, economic costs, foodborne disease costs, *Salmonella* control, salmonellosis, 936
Roasted turkey, consumer preference, fatty material injection, turkey broth injection, 160*
Roaster, ascites, broiler, environmental temperature, 87*
Roaster weight, breed effect, nutrition, temperature regimen, 96*
Rocket electrophoresis, bovine serum albumin injection, host immunity, northern fowl mite, two-dimensional immunoelectrophoresis, 1015
Roost temperature, conductive heat loss, heat stress, metabolic rate, panting, 124*
Rooster, spermatozoal integrity, zinc, 103*
Rooster-intact, cecectomized or cannulated, diaminopimelic acid, lysine, microorganisms, 1345†
Rous sarcoma virus, anti-tumor response, *B* complex genotype, histocompatibility complex, 164*
B complex genotype, tumor regression, 1491
genetic lines, immunity, intracerebral inoculation, neurological behavior, 1123

Roxarsone, bacitracin, body weight, broiler, narasin, 103
broiler chick, glutathione peroxidase, selenium, 41*
Rye-ryegrass pasture, beef calves performance, broiler litter-corn diet, 29*
Salinomycin, anticoccidial drugs, maduramicin, narasin, nicarbazin, 35*
body weight, turkey poult, viability, vomitoxin, 149†
broiler performance, methionine, monensin, nicarbazin, 33*
Salmonella, antibiotic resistance, broiler carcass, market survey, 55*
Escherichia coli, food chain, kanamycin, 699
kanamycin, transconjugant, 689
bactericide, lactic acid, poultry-processing water, 69*
broiler, chemical additives, microbial contamination, poultry-processing water, 10*
broiler carcass, chemical preservative, feed, Myco Curb®, 1225†
detection, DNA probe, poultry-processing water, 80*
broiler chick, colonization dose-50%, 71*
broiler meat, microwave cooking, 71*
carbon sources, coliforms, fermentation, manure-cage layer, 74*
chicken meat, high energy electrons, irradiation pasteurization, 165*
irradiation, poultry products, sensory quality, shelf life, 36*
Salmonella control, *Salmonella* control economics, 920
***Salmonella* attachment**, ceca, mucin layer, 26*
***Salmonella* colonization control**, broiler growout, broiler processing, competitive exclusion, 928
***Salmonella* contamination in poultry**, Food Safety and Inspection Service, poultry inspection, *Salmonella* control, 944
***Salmonella* control**, broiler processing, chilling, evisceration, scalding, 933
competitive exclusion, epidemiological evaluation, 1026
economic costs, foodborne disease costs, risk assessment, salmonellosis, 936
Food Safety and Inspection Service, poultry inspection, *Salmonella* contamination in poultry, 944
Salmonella, *Salmonella* control economics, 920
***Salmonella* control economics**, *Salmonella*, *Salmonella* control, 920
***Salmonella* isolation**, broiler carcass, ELISA, feed, gene probe, 921
Salmonella typhimurium, broiler chick, detection, enteropathogen, 52*
chick, intestine, mannose sensitivity, receptors, 133*
Salmonellosis, chick, cyclosporine, immunosuppression, 176*
economic costs, foodborne disease costs, risk assessment, *Salmonella* control, 936
Salt, egg yolk, physical manipulation, shaking, texture, 1264
Salt concentration, broiler breast tissue, fiber orientation, 122*
Salt soluble protein, exudate, marinade composition, tumbling, turkey meat, 162*
Salted yolk paste, egg yolk, food-poisoning bacteria, lecithovitellin reaction, *Staphylococcus aureus*, 261
Sand, confounding experimental results, energy supplementation, growth assay, oil, 169*
Sanitizers, fertility, hatchability, turkey egg, 144*
Sarcomere length, broiler, muscle tenderness, pH, postmortem temperature, 54*
broiler breast muscle, R value, rigor development, 635
Satellite cell, muscle growth, myogenic, turkey, 121*
Sausage, fat sources, fatty acid profile, lipid oxidation, 91*
Scalding, broiler processing, chilling, evisceration, *Salmonella* control, 933
Scalding temperature, broiler, feather release forces, scalding time, stunning *vs.* nonstunning, 1069
Scalding time, broiler, feather release forces, scalding temperature, stunning *vs.* nonstunning, 1069
Scalding-spay scalding, bacterial contamination, broiler carcass, cross-contamination, 77*
Scan sheet scoring, computerized contest scoring, judging contest, 140*
Scientific aspect of poultry industry, business aspect of poultry industry, curriculum, poultry programs of the future, student recruitment, 879
SDS-PAGE analysis, *Mycoplasma gallisepticum*, *Mycoplasma* species, 116*
Season of the year, age, body weight, broiler male, sudden death syndrome, 1243
Seasonal change, Japan, layer performance year round, open-sided houses, photoperiod, 1130
Secondary plate development, embryogenesis, morphogenesis, pigeon, 865
Secretion rate, age, growth hormone, metabolic clearance rate, protein restriction, 120
Secretory cell, bursa, dendritic cell, 16*
Seed-processing methods, canola seed, full-fat canola seed, pelleting, turkey, 1731
SEHCAT, bile acids, broiler chick, intestine, liver, 17*
Selectable marker, gene transfer, retrovirus vector, 42*
Selection, abdominal fat, body weight, broiler sire strain, feed conversion, 10
broiler, 65*
broiler breeder, dietary energy, quantitative traits, 6*
genetic correlation, sexual dimorphism, turkey, 166*
growth, nutritional requirements, quail-Japanese, 22*
Selection differentials, growth rate, heritability, protein and energy, quail-Japanese, 519
Selection index, egg mass, egg production, genetic parameters, shell strength, 42*
Selection lines, antibody production, environmental temperature, immune response, sheep red blood cells, 79*
heat stress, immune response, sheep red blood cells, 79*
Selection response, adult selection, genetic merit, juvenile selection, layer performance, 1655
Selenium, bioavailability, broiler male chick, time, tissue uptake, 1295
chick, tissue uptake, 1585
broiler chick, glutathione peroxidase, glutathione reductase, riboflavin, 149*
roxarsone, 41*
Selenium deficiency, macrophages, vitamin E deficiency, 141*
Self-select diets, calcium intake, feed intake, layer, 24*
energy level, protein level, pullet, sexual maturity, 153*
lighting program, pullet, sexual maturity, temperature, 152*
Semen, blood plasma, seminal plasma, testosterone, turkey breeder, 1461
broiler breeder evaluation, glycerol, tissue culture, 23*
fertility, freezing, spermatozoal organelles, ultrastructure, 819
Iraqi chickens, *Turella* yeast, 46*
Semen characteristics, body composition, broiler breeder male, feeding regimen, 835

Semen diluent, antibiotics, sperm lysis, sperm motility, turkey, 131
Semen extender, artificial insemination, fertility, spermatozoa-damaged, turkey semen, 1483†
carboxymethyl cellulose, fertility, semen viscosity, 1218
Semen production, body composition, broiler breeder, feed restriction, 34*
body weight, broiler breeder male, testes morphology, testis weight, 660
chick, precocious puberty, tamoxifen, 1642†
egg production, feed conversion, fertility and hatchability, rabbit excreta, 168*
ejaculate, spermatogenesis, turkey, 327
reproductive performance, thyroidectomy, turkey, 110*
Semen viscosity, carboxymethyl cellulose, fertility, semen extender, 1218
Semen yield, sperm concentration, turkey breeder growth, water restriction, 1236†
Semi-isolation brooding, broiler chick performance, management, weight gain, 96*
Seminal plasma, blood plasma, semen, testosterone, turkey breeder, 1461
Sensory evaluation, chicken patties, reheating methods, storage time, thiobarbituric acid values, 736
Sensory profiling, chicken patties, off-flavor, reheating methods, thiobarbituric acid, 112*
Sensory properties, irradiation, sodium reduction, turkey frankfurters, 1797†
Sensory quality, breast up *vs.* breast down cooking, canola meal, cooking procedure, turkey, 680†
irradiation, poultry products, *Salmonella*, shelf life, 36*
Serotonin, blood leukocyte, dopamine, migration, norepinephrine, 116*
bovine serum albumin, chick, dopamine, ELISA, 100*
electric stimulation, prolactin, turkey, 93*
Serum, carotenoid composition, egg, lutein, yolk, 608
cholesterol, egg, full-fat canola seed, omega-3 fatty acids, 129*
melatonin level, pineal gland, *rc* chickens, retina, 66*
Sesbania seed, broiler chick performance, layer performance, 11*
Sex chromosome, barring, dermal melanin, linkage relationship, recessive white skin, 530
Sex separate rearing, amino acid(s), broiler, energy level, 39*
Sex-linkage, avian leukosis virus, congenital transmission, endogenous virus, slow-feathering, 1668
Sex-linked coloration, chick, feathers, fiber optics, 1680
Sex-linked gene, genetics, *Pavo cristatus*, peafowl, plumage color mutation, 158*
Sexual dimorphism, genetic correlation, selection, turkey, 166*
Sexual dysfunction, hypothalamus, lesions, obesity, 157*
Sexual maturity, body weight, dwarf Leghorns, reproduction, 143*
broiler breeder, egg production, feed restriction, 1001
egg production, propylthiouracil, pullet, 107*
energy level, protein level, pullet, self-select diets, 153*
lighting program, pullet, self-select diets, temperature, 152*
Shaking, egg yolk, physical manipulation, salt, texture, 1264
Shank length, body density, divergent selection, quail-Japanese, 88*
Shank width, body weight, leg bones, leg muscles, turkey, 1660
Shear force, breast muscle, marination, moisture binding, rigor, 175*
Shear strength, feed and water withdrawal, gastrointestinal tract, moisture content, water withdrawal, 845†
Sheep red blood cells, anamnestic response, antibody response, 855†
antibody production, environmental temperature, immune response, selection lines, 79*
heat stress, immune response, selection lines, 79*
bursal cell transfer, dopamine, immune response, plaque-forming cell transfer, 88*
Shelf life, irradiation, poultry products, *Salmonella*, sensory quality, 36*
Shell calcium, egg weight, shell thickness, shell weight, specific gravity, 159*
Shell gland, egg shell strength, molt-induced, phospholipids, 427
fat, lecithin, phospholipids, shell strength, 8*
Shell quality, acid-base balance, gypsum, layer diet, pulverized and particle size limestone, 20*
ahemeral photoperiod, egg production, laying photoperiod, rearing photoperiod, 391
blood levels, calcium, layer, phosphorus, 68*
broiler breeder, feeding time, hatchability, slow *vs.* fast feathering, 538
calcium source, egg production, limestone solubility, particle size, 66*
chloride, ion balance, layer diet, sodium, 39*
defective eggs, genetic associations, oviposition rate, shell-less egg, 123*
economic impact, egg breakage, uncollectable eggs, 1801†
Shell quality measurements, egg breakage, egg-processing plant, 162*
Shell strength, calcitriol, layer, 65*
egg mass, egg production, genetic parameters, selection index, 42*
fat, lecithin, phospholipids, shell gland, 8*
Shell thickness, egg weight, shell calcium, shell weight, specific gravity, 159*
Shell treatment, age of bird, egg quality, egg storage position, yolk position, 75*
Shell weight, egg weight, shell calcium, shell thickness, specific gravity, 159*
Shell-less culture, chick embryo, mammalian embryo, 55*
Shell-less egg, amino acid analysis, egg shell membranes, egg shell quality, soft-shelled egg, 1203
defective eggs, genetic associations, oviposition rate, shell quality, 123*
Shortening quality, chicken patties, frying conditions, lipid content, moisture content, 9*
Showmanship guidelines, 4-H, duck, fitting guidelines, turkey, 167*
Silicon, aluminum, broiler, tibial dyschondroplasia, 84*
Simulation model, broiler grower, interregional competition, profitability, 89*
Simulation program, broiler growout, computer program, optimization, temperature, 534
Skeletal diseases, broiler performance, feed restriction, lighting program, 68*
Skeletal measurments, body weight, genetic correlation, heritability, turkey, 1378
Skeletal muscle, histochemistry, muscle fiber type, 973
Skeletal muscle cell culture, chick embryo, leupeptin, proteolysis, 87*
Skin strength, broiler crosses, feather removal, feeding regimen, kosher processing, 125*
Slaughter yield, abdominal fat, broiler sire strain, feed conversion, weight gain, 16
Slow *vs.* fast feathering, broiler breeder, feeding time,

hatchability, shell quality, 538
Slow-feathering, avian leukosis virus, congenital transmission, endogenous virus, sex-linkage, 1668
tumors, 1674
Small intestine, antibiotics, broiler growth, 43*
Small scale plants, broiler processing, budgeting techniques, costs, 76*
Sodium, acid-base balance, dietary calcium, dietary chloride, egg shell quality, 750
acid-base status, broiler growth, sodium bicarbonate, 151*
chloride, ion balance, layer diet, shell quality, 39*
Sodium aluminosilicate, egg shell quality, egg specific gravity, zeolite, 577
layer performance, phosphorus, 31*
Sodium availability, broiler chick, Ethacal® feed component, 15*
Ethacal® feed component, layer, 120*
Sodium bicarbonate, acid-base status, broiler growth, sodium, 151*
Sodium pyrophosphate, carcass size, marinated broiler, moisture absorption, moisture retention, 510†
Sodium reduction, irradiation, sensory properties, turkey frankfurters, 1797†
Sodium salts, dark turkey meat, protein, storage, 103*
storage time, water added, 134*
Sodium sulfate, broiler performance, methionine level, 159*
Sodium vapor, broiler growth performance, fluorescent, incandescent, light source, 43
Soft-shelled egg, amino acid analysis, egg shell membranes, egg shell quality, shell-less egg, 1203
Solubility, calcium source, layer digestive system, limestone, particle size, 142*
limestone, particle size, 29*
Somatotrophs, adenohypophysis, age, chick, growth hormone, 487
Sorghum, amino acid(s), crude protein, regression equations, 40*
broiler, corn, fatty acids, processing yield, 130*
Soy protein, broiler chick, hypersensitivity, immune response, 104*
Soybean meal, apparent metabolizable energy, corn, true metabolizable energy, 10*
biotin bioavailability, canola meal, cereal grains, turkey poult, 1274
body weight, broiler chick, feed conversion, tibial dyschondroplasia, 21*
pancreas weight, turkey, urease index, 15*
broiler chick, protein solubility, urease activity, 11*
broiler performance, cottonseed meal, 76*
corn-soy-gelatin basal diet, methionine activity, methionine assay, 88
digestibility, oligosaccharides, transit time, 109*
egg mass, egg production, layer, lysine assay, 49*
protein, toasting temperature, trypsin inhibitor, urease index, 20*
Soybeans, broiler diet, trypsin inhibitor, 37*
Soybeans roasted *vs.* raw, broiler starter diet, soybeans-extruded, turkey starter diet, 163*
Soybeans-extruded, broiler starter diet, soybeans roasted *vs.* raw, turkey starter diet, 163*
Specific gravity, broiler breeder performance, Ethacal® feed component, 35*
egg weight, shell calcium, shell thickness, shell weight, 159*
Sperm concentration, semen yield, turkey breeder growth, water restriction, 1236†
Sperm lysis, antibiotics, semen diluent, sperm motility, turkey, 131
Sperm motility, antibiotics, semen diluent, sperm lysis, turkey, 131
Sperm viability, broiler breeder, divergent selection, fluorometry, 55*
Spermatogenesis, ejaculate, semen production, turkey, 327
Spermatozoa-chicken, cell culture, glycerol, 135
Spermatozoa-damaged, artificial insemination, fertility, semen extender, turkey semen, 1483†
Spermatozoal integrity, rooster, zinc, 103*
Spermatozoal organelles, fertility, freezing, semen, ultrastructure, 819†
Spleen, bursa, cyclophosphamide, dendritic cell, 114*
Splenomegaly, adrenocorticotropin, ascorbic acid, cell-mediated immunity, graft *vs.* host reaction, 313
Spoilage, dimethyl sulfide, egg products, odor, uracil, 123*
Sporozoites, *Eimeria*, foreign hosts, 49*
Spray scalding, broiler, feather release, microbiological contamination, picking feathers, 12*
Spray-dried egg white, cooked gels, mechanically deboned meat, water-washed, 73*
Standard *vs.* all-corn diet, broiler breeder male, diet, nitrogen retention, 144*
Staphylococcus aureus, egg yolk, food-poisoning bacteria, lecithovitellin reaction, salted yolk paste, 261
genetic analysis, wattle reaction, 70*
Sterigmatocystin, blood chemistry, chick, metabolites, organ weight, 502
Steroid hormones, epidermal growth factor, gonadotrophins, granulosa cell, *in vitro*, 814
hen plasma, 167*
Steroidogenesis, antibiotics, bovine luteinizing hormone, ovarian tissue, 1486†
layer, luteinizing hormone, ovarian follicle, 145*
ovarian cells, prolactin, turkey, 95*
Stocking density, litter moisture, niacinamide, turkey poult performance, 24*
Storage, dark turkey meat, protein, sodium salts, 103*
Storage condition, egg, internal quality, 133*
Storage container, fertility, frozen semen, 333†
Storage time, bacterial flora, broiler meat, oxidative stability, 2*
chicken patties, reheating methods, sensory evaluation, thiobarbituric acid values, 736
dark turkey meat, sodium salts, water added, 134*
functional properties of myofibrils, lipid oxidation, myofibrils, water activity, 128*
Strain, age, egg yolk, layer, 7*
Streptococcus faecium, body weight, broiler, probiotic, 132*
Stress, ascorbic acid level, turkey neonate, 134*
beak trimming, broiler chick performance, feed efficiency, heterophil:lymphocyte ratio, 166*
behavior, chick performance, electric shock-experimental system, health, 1791†
broiler breeder, corticosterone, feed restriction, tryptophan, 119*
corticosterone, heritability, quail-Japanese, 25
prostaglandin-E_2, quail-Japanese, 32*
Structural unit, calcium reserve, egg shell, 77*
Student performance, majors *vs.* nonmajors, poultry component in introductory course, 149*
Student recruitment, business aspect of poultry industry, curriculum, poultry programs of the future, scientific aspect of poultry industry, 879
FFA and 4-H, poultry majors, poultry programs of the future, youth programs, 886

Stunning *vs.* nonstunning, broiler, feather release forces, scalding temperature, scalding time, 1069
Sublines, fertility, hatchability, immunogenetics, inbreeding, 859
Substrate specifity, amino acid(s), D-amino acid oxidase, duckling, 851†
Sudden death syndrome, age, body weight, broiler male, season of the year, 1243
broiler, feed efficiency, feed restriction, growth rate, 1102†
broiler male, estradiol, 156†
Sulfur amino acid level, 35 S-cystine, 35 S-methionine, egg albumen, 78*
Sulfur amino acid requirement, age, layer, methionine requirement, 170*
Sulfur amino acids, layer requirements, lysine, 130*
Sunflower meal, broiler, feed efficiency, pelleting, 175*
Sunflower seeds, Aureomycin, egg production, pelleting, pullet growth, 445
Supplemental equipment, broiler chick performance, feed, water, 124*
Surgery stress, arginine vasotocin, broiler, heat stress, mesotocin, 171*
Surimi-wash process, gel-strength, mechanically deboned meat, poultry meat color, 91*
Synergy, aflatoxin, broiler male chick, mycotoxin, T-2 toxin, 1418
T cell-mediated response, antibody response, histocompatibility complex, immunocompetence, phagocytosis, 989
T-2 toxin, aflatoxin, broiler male chick, mycotoxin, synergy, 1418
broiler, chlortetracycline, gentian violet, 99*
T_3,T_4, broiler pullet, growth hormone, insulin, pulsatile, 168*
T_3,T_4 doses, antibody production, growth hormone, leukocyte distribution, lymphoid organs, 92*
B-cells, growth hormone, natural killer cells, T-cells, 93*
Table egg production, broiler production, commercial chicken industry, educational tool, video, 134*
educational tool, video, 46*
Tamoxifen, chick, precocious puberty, semen production, 1642†
Tannin content, broiler performance, fat, grain sorghum, metabolizable energy, 80*
T-butyl hydroperoxide, erythrocytes, hemolysis, plasma, uric acid, 1632
T-cells, *B*-cells, growth hormone, natural killer cells, T_3,T_4 doses, 93*
Teaching poultry, extension program, poultry programs of the future, poultry students, 878
Teaching tool quiz, educational poultry display, 63*
Temperature, body weight, feed restriction, molt, 137*
broiler, chilling time, glycol solution, packaged carcasses, 154*
broiler growout, computer program, optimization, simulation program, 534
carcass composition, energy, layer, metabolizable energy utilization, 136*
egg type pullet, mash, pellets *vs.* mash, weight gain, 375
environmental stress, layer performance, relative humidity, vitamin C, 67*
fertility, naked neck, reproductive traits, 176*
lighting program, pullet, self-select diets, sexual maturity, 152*
molt, postmolt performance, 126*
Temperature environment, body weight gain, broiler performance, Ethacal® feed component, 18*
Temperature regimen, breed effect, nutrition, roaster weight, 96*
Temperature-hot weather, egg production, energy level, layer diet, 108*
Tenderness, accelerated rigor, electric stimulation, minimum time process, nonaged tenderness, 40*
adenosine triphosphate, electric stimulation, pH, poultry carcass, 106*
Allo-Kramer shear, broiler breast meat, texture profile analysis, 35*
boning time, broiler breast fillets, electric stimulation, post-mortem metabolism, 150*
brine chilling, broiler breast fillets, cooking method, 715
brine solution *vs.* water, broiler breast fillets, chilling temperature, hot boning, 1545
broiler breast meat, textural quality, wing restraint, 275
Teratogenicity, 3,3′,4,4′-tetrachlorobiphenyl, duck embryo, goose embryo, herring gull embryo, 52
Testes morphology, body weight, broiler breeder male, semen production, testis weight, 660
Testicular development, feather color, photoperiod, quail-Japanese, 1098†
Testis, electron microscopy, epididymis, gossypol acetic acid, quail-Japanese, 961
Testis weight, body weight, broiler breeder male, semen production, testes morphology, 660
Testosterone, blood plasma, semen, seminal plasma, turkey breeder, 1461
corticosterone, feed restriction, quail-Japanese, 121*
Textural quality, broiler breast meat, tenderness, wing restraint, 275
Texture, Allo-Kramer shear, broiler breast meat, texture profile analysis, 1549
broiler breast meat, electric stimulation, hot boning, poultry meat color, 1536
egg yolk, physical manipulation, salt, shaking, 1264
Texture profile, broiler breast muscle, cooking method, deboning time, post-mortem aging, 23*
electric stimulation, voltage levels, 23*
Texture profile analysis, Allo-Kramer shear, broiler breast meat, tenderness, 35*
texture, 1549
Theca cell, 17α-hydroxylase, hypophysectomy, ovarian follicle, quail-Japanese, 1636
Thermal perch, conductive heat loss, foot temperature, heat stress, layer, 142*
Thermoregulation, acid-base status, blood, heat stress, layer, 113*
Thiamin content, breast muscle, cooking method, purification variables, 72
Thiamin deficiency, anorexia, energy metabolism, feed intake, turkey, 143*
Thigh, breast muscle, broiler, exercise patterns, muscle growth, 1708
Thigh muscle, breast muscle, processing treatment, turkey Canadian bacon, 162*
Thiobarbituric acid, chicken patties, off-flavor, reheating methods, sensory profiling, 112*
Thiobarbituric acid values, chicken patties, reheating methods, sensory evaluation, storage time, 736
Thiouracil, dietary protein, plasma thyroxine, quail-Japanese, 28*
egg weight loss, embryonic development, embryonic mortality, quail-Japanese, 28*
Thiourea, hatch time, house fly, larvicide, manure, 407

Threonine, body weight, lysine, turkey poult, valine, 140*
Threonine requirement, amino acid(s), body weight, broiler male chick, feed conversion, 108
Thymidine uptake, antibody production, immune response, light regimen, white blood cell, 114*
Thyroid dysfunction, acid water, broiler breeder, 109*
Thyroidectomy, reproductive performance, semen production, turkey, 110*
Thyroxine, egg production, molt, photorefractoriness, turkey, 21*
Tibial dyschondroplasia, acid-base balance, broiler chick, mineral anions, 148*
aluminum, broiler, silicon, 84*
body weight, broiler chick, feed conversion, soybean meal, 21*
broiler chick, epiphyseal growth plate, growth hormone, 82*
vitamin A, zeolite, 113
calcium, chloride, phosphorus, zeolite, 1436
Time, bioavailability, broiler male chick, selenium, tissue uptake, 1295
Tissue culture, broiler breeder evaluation, glycerol, semen, 23*
Tissue distribution, egg, ochratoxin A, quail-Japanese, whole body autoradiography, 707
Tissue tropism, infectious bronchitis virus, pathogenesis, 60*
Tissue uptake, bioavailability, broiler male chick, selenium, time, 1295
chick, selenium, 1585
Tissue-specific expression, chicken embryo, gene transfer, retrovirus vector, 34*
chloramphenicol acetyltransferase, gene transfer, retrovirus vector, 156*
Toasting temperature, protein, soybean meal, trypsin inhibitor, urease index, 20*
Toe ash, bioavailability, body weight, phosphorus, turkey, 96
Toltrazuril, anticoccidial efficacy, BAY Vi 9142, coccidiosis, postinfection, 122*
Toxicity, butylated hydroxyanisole, cysteine, lathyrism, quail-Japanese chick, 153†
Trachea, aorta, *Bordetella avium*, elastin, turkey poult, 174*
***Trans*/18:1**, biotin, chick, fatty acids, liver lipids, 45*
***Trans*/*cis* 18:1**, blended feed-grade fats, fatty acid composition, gas-liquid chromatography, 307
chick tissue, fatty acids, hydrogenated, 778
Transconjugant, antibiotic resistance, *Escherichia coli*, kanamycin, *Salmonella*, 689
Transit time, digestibility, oligosaccharides, soybean meal, 109*
Transsulfuration enzyme activity, cystine, layer performance, methionine, 78*
Triticale, body weight, broiler diet, litter condition, 1743
dietary protein, energy, reproductive performance, turkey semen, 1619
True amino acid availability, amino acid(s), bioavailability, *in vitro* digestion, plant feedstuffs, 78
True metabolizable energy, apparent metabolizable energy, corn, soybean meal, 10*
meat and bone meal, metabolizable energy, 280
bioassay, excreta collection, glucose, water, 1192
Trypsin inhibitor, broiler diet, soybeans, 37*
protein, soybean meal, toasting temperature, urease index, 20*
Tryptophan, aggressive behavior, broiler breeder male, feed restriction, 155*
Bordetella avium, cilia, collagen, turkey poult, 42*
broiler breeder, corticosterone, feed restriction, stress, 119*
niacin, trytophan:niacin conversion, turkey poult, 148*
Tryptophan requirement, amino acid(s), broiler male chick, 1174
Trytophan:niacin conversion, niacin, tryptophan, turkey poult, 148*
Tumbling, exudate, marinade composition, salt soluble protein, turkey meat, 162*
Tumor regression, *B* complex genotype, Rous sarcoma virus, 1491
Tumors, avian leukosis virus, congenital transmission, endogenous virus, slow-feathering, 1674
***Turella* yeast**, Iraqi chickens, semen, 46*
Turf Bermudagrass, carotene, drying method, grass-clipping length, xanthophyll, 1644†
Turkey, 4-H, duck, fitting guidelines, showmanship guidelines, 167*
5-azacytidine, DNA methylation, gene expression, prolactin, 1794†
methylation, prolactin, 86*
abdominal fat pad, cloaca caliper measurment, 29*
abnormal eggs, cage *vs.* floor housing, embryonic mortality, fertility, 1455
adrenocortical cell subpopulation, corticosteroidogenesis, 105*
amino acid(s), fat, protein, 4*
anorexia, energy metabolism, feed intake, thiamin deficiency, 143*
antibiotics, semen diluent, sperm lysis, sperm motility, 131
antibody response, egg production, Newcastle disease virus, *Pasteurella multocida*, 1372
Pasteurella multocida, 149*
bacitracin, body weight, lasalocid, 99*
bioavailability, body weight, phosphorus, toe ash, 96
body weight, carcass composition, genetic correlation, heritability, 1388
carcass yield, parts composition, 153*
genetic correlation, heritability, skeletal measurements, 1378
leg bones, leg muscles, shank width, 1660
pancreas weight, soybean meal, urease index, 15*
breast up *vs.* breast down cooking, canola meal, cooking procedure, sensory quality, 680†
broiler, economics, layer, naked oats pricing, 10*
broody, egg production, egg quality, progesterone, 72*
broody hen management, prolactin, 128*
canola seed, full-fat canola seed, pelleting, seed-processing methods, 1731
carcass fat, dietary fat, feed efficiency, growth rate, 420
carcass quality, compensatory growth, organ proportions, protein undernutrition, 1760
catecholamines, chicken, feed intake, peptides, 74*
conductance, egg shell permeability, hatchability, 9*
dietary fat, hepatic tocopherol concentration, liver, vitamin E, 1814†
disease challenge, egg production, *Pasteurella multocida*, 1807†
egg production, molt, photorefractoriness, thyroxine, 21*
ejaculate, semen production, spermatogenesis, 327
electric stimulation, prolactin, serotonin, 93*
estradiol, ovarian growth, pregnant mare serum gonadotropin, prolactin, 319
follicle-stimulating hormone, radioreceptor assay, 95*
fowl cholera, KSCN extract, *Pasteurella multocida*, vac-

cination, 26*
genetic correlation, selection, sexual dimorphism, 166*
glucagon, insulin, lame *vs.* normal, leg abnormalities, 97*
glycerol trioleate, pancreatic lipase, 85*
glycoprotein hormone, hybridoma, luteinizing hormone, monoclonal antibodies, 117*
heparin, nonesterified fatty acids, very low density lipoprotein, 51*
hypothalamus, photostimulation, prolactin, 138*
muscle growth, myogenic, satellite cell, 121*
ovarian cells, prolactin, steroidogenesis, 95*
peptide histidine isoleucine, prolactin-releasing factor, 141*
reproductive performance, semen production, thyroidectomy, 110*

Turkey breast meat, heme-complex, nicotinamide, poultry meat color, pyridine, 44*

Turkey breeder, bioelectrical impedance analysis, body composition, fat free mass, 92*
blood plasma, semen, seminal plasma, testosterone, 1461
nutrition, progeny performance, 170*

Turkey breeder growth, semen yield, sperm concentration, water restriction, 1236†

Turkey breeder performance, body composition, energy, protein, 17*
corn-soybean meal diet, fertility, hatchability, niacin supplementation, 336†
light regimen, 173*

Turkey broth injection, consumer preference, fatty material injection, roasted turkey, 160*

Turkey Canadian bacon, breast muscle, processing treatment, thigh muscle, 162*

Turkey egg, biotin level, fertility, hatchability, 30*
egg shell permeability, hatchability, iodine, 67*
fertility, hatchability, sanitizers, 144*

Turkey egg hatchability, embryonic mortality, high altitude, incubation temperature, oxygen tension, 956

Turkey embryo, antibiotic treatment, glucose supplement, hatching metabolism, 463

Turkey fertility, anti-sperm antibodies, ELISA, immunohistochemistry, 20*

Turkey frankfurters, irradiation, sensory properties, sodium reduction, 1797†

Turkey genotypes, growth, muscle development, 22*

Turkey growout, computer spreadsheet templates, contract, financial attractiveness, 142*

Turkey growth, dehulling, meat flavor, white lupine, 596

Turkey growth rate, genetic lines, growth hormone, 826

Turkey hen, added fat in diet, quaifenesin in diet, reproductive performance, 114*

Turkey liver explants, buffers, glycogenolysis, hormones, 147*

Turkey male, body weight, dietary fat, feed efficiency, protein, 1281
carcass composition, compensatory growth, feed restriction, 1407

Turkey male performance, body weight, feed efficiency, feed intake, protein restriction, 130*

Turkey management, feed form, floor space, lighting environment, 127*

Turkey meat, exudate, marinade composition, salt soluble protein, tumbling, 162*

Turkey meat product form, futher processing, microbiological growth support, nutrient availability, particle size, 1269

Turkey muscle, delayed glycolysis, electric stunning, muscle pH, protein solubility, 1062
growth, myopathies, 173*

Turkey neonate, ascorbic acid level, stress, 134*

Turkey performance, barley, corn texture, grit, pellets, 101*
barley brewers grain meal, broiler performance, cholesterol, 175*
body composition, compensatory growth, protein undernutrition, 1750
body weight, virginiamycin, 94*
carcass composition, carcass yield, corn gluten feed, feed efficiency, 585
color-filtered light, growth, photoperiod, 1505
endocrine glands, light environment, light intensity, population density, 1518
floor type, litter condition, 66*

Turkey poult, 17β-estradiol dipropionate, lipase activity, pancreas, 84*
age, diet, intestinal disaccharidases, 153*
aorta, *Bordetella avium*, elastin, trachea, 174*
biogenic amines, duodenum, hypothalamus, mortality, 13*
biotin bioavailability, canola meal, cereal grains, soybean meal, 1274
body weight, feed conversion, leg abnormalities, niacin requirement, 760
lysine, threonine, valine, 140*
salinomycin, viability, vomitoxin, 149†
Bordetella avium, cilia, collagen, tryptophan, 42*
calcium requirement, phosphorus requirement, rickets, 150*
choline, methionine, 9*
glucose, glycogen, intestine, yolk sac, 493
lipase activity, pancreas, yolk membrane, 1089
litter, rickets, 106*
niacin, tryptophan, trytophan:niacin conversion, 148*
riboflavin requirement, 32*

Turkey poult performance, litter moisture, niacinamide, stocking density, 24*

Turkey roast, cooking, meat composition, meat source, 1810†

Turkey roast size, cooking time, oven type, 139*

Turkey sausage, beef sausage, fat content, meat flavor, quality attributes, 1075

Turkey semen, artificial insemination, fertility, semen extender, spermatozoa-damaged, 1483†
dietary protein, energy, reproductive performance, triticale, 1619

Turkey starter diet, broiler starter diet, soybeans roasted *vs.* raw, soybeans-extruded, 163*

Turkey tom, meat yield, 43*

Turkey tom performance, amino acid(s), growth rate, lysine, methionine, 86*
energy, pellets, protein level sequence, 170*

Two-dimensional immunoelectrophoresis, bovine serum albumin injection, host immunity, northern fowl mite, rocket electrophoresis, 1015

Two-slope broken line method, lysine requirement, protein source, 53*

Ulcerative enteritis, antimicrobial agents susceptibility, broiler, *Clostridium colinum*, 1424
Clostridium, psittacine, 75*

Ultrastructure, fertility, freezing, semen, spermatozoal organelles, 819

Uncollectable eggs, economic impact, egg breakage, shell quality, 1801†

Undergraduate student, poultry science, recruitment methods, 160*

Uracil, dimethyl sulfide, egg products, odor, spoilage, 123*
Urea, blood chemistry, formaldehyde, layer, pellet binder, 37*
Urease activity, broiler chick, protein solubility, soybean meal, 11*
Urease index, body weight, pancreas weight, soybean meal, turkey, 15*
protein, soybean meal, toasting temperature, trypsin inhibitor, 20*
Uric acid, erythrocytes, hemolysis, plasma, t-butyl hydroperoxide, 1632
feeding time, Leghorn male, meal feeding, plasma, 345†
Urinary excretion, broiler chick, DL-methionine, DL-methionine hydroxy analog, L-methionine, 14*
Urolithiasis, acid-base balance, calcium level, pullet, 90*
dietary acidification, dietary alkalinization, layer performance, renal function, 1694
dietary calcium, infectious bronchitis virus, kidney function, pullet chick, 1250
Uterus, myoelectrical activity, oviposition, vagina, 1615
Vaccination, embryo, hatching egg injection machine, incubation, Marek's disease, 110*
fowl cholera, KSCN extract, *Pasteurella multocida*, turkey, 26*
Vaccine, coccidiosis, *Eimeria tenella*, recombinant DNA, 72*
Vagina, myoelectrical activity, oviposition, uterus, 1615
Valgus-varus, biotin deficiency, bone modeling, broiler chick, linoleate, 590
Valine, body weight, lysine, threonine, turkey poult, 140*
broiler chick, isoleucine, leg abnormalities, leucine, 85*
Variances, fasted rooster, feather meal, metabolizable energy assays, 1188
Varus-angular bone deformity, quantitative evaluation method, 1647†
Ventilation, broiler, evaporative cooling, fan cooling, pad cooling, 98*
Very low density lipoprotein, chick embryo, cholesterol, growth hormone, insulin-like growth factor-1, 71*
heparin, nonesterified fatty acids, turkey, 51*
Viability, body weight, salinomycin, turkey poult, vomitoxin, 149†
Vicine, egg production, egg quality, field beans, layer performance, 455
Video, broiler production, commercial chicken industry, educational tool, table egg production, 134*
educational tool, table egg production, 46*
Villafranquina chickens, black-tailed red pattern, dark brown gene, gene interactions, plumage color, 351
Viral diagnosis, chick, infectious bronchitis virus, recombinant DNA, 69*
Virginiamycin, body weight, turkey performance, 94*
broiler yield, dietary energy, 1222
Virus isolation, immunity, infectious bronchitis virus, 174*
Viscosity, barley, chick performance, enzyme supplementation, weight gain, 147*
Vitamin A, bone development, egg injection, embryo, vitamin D_3, 58*
broiler chick, tibial dyschondroplasia, zeolite, 113
Vitamin C, broiler, processing yield, 85*
environmental stress, layer performance, relative humidity, temperature, 67*
Vitamin D, 1,25-dihydroxyvitamin D_3, egg shell quality, layer performance, 166*
aluminum, broiler chick, phosphorus metabolism, 19*
Vitamin D_3, bone development, egg injection, embryo, vitamin A, 58*
Vitamin E, broiler meat quality, carcass weepage, lipid stability, 38*
dietary fat, hepatic tocopherol concentration, liver, turkey, 1814†
Vitamin E deficiency, macrophages, selenium deficiency, 141*
Vitelline membrane, egg, glycoprotein, membrane layers, ovomucin, 476
Vitellogenin gene, DNA modifications, egg yolk protein precursor, quail-Japanese, 70*
Volatile analysis, cooking method, marinated chicken fillet, moisture content, yield, 128*
Voltage levels, broiler breast muscle, electric stimulation, texture profile, 23*
Vomitoxin, body weight, salinomycin, turkey poult, viability, 149†
Washing, composition, functional properties, mechanically deboned meat, 87*
Wastewater treatment, BOD removal, COD removal, hydrogen peroxide, poultry-processing water, 24*
Water, bioassay, excreta collection, glucose, true metabolizable energy, 1192
broiler chick performance, feed, supplemental equipment, 124*
Water activity, functional properties of myofibrils, lipid oxidation, myofibrils, storage time, 128*
Water added, dark turkey meat, sodium salts, storage time, 134*
Water conservation, flow volume measurement, poultry-processing plant, 119*
Water consumption, broiler, prediction equation, water:feed ratio, 131*
Water intake, B-endorphin, body temperature, feed intake, 25*
broiler, cellulose, crude fiber, 169*
feed intake, heat stress, 38*
broiler performance, cyclic watering, feed intake, 378
Water quality, broiler performance, chlorine, waterers, 125*
Water restriction, semen yield, sperm concentration, turkey breeder growth, 1236†
Water withdrawal, feed and water withdrawal, gastrointestinal tract, moisture content, shear strength, 845†
Water-washed, cooked gels, mechanically deboned meat, spray-dried egg white, 73*
Water:feed ratio, broiler, prediction equation, water consumption, 131*
Waterer location, cage shape, layer performance, nipple waterer, waterers per cage, 89*
Waterers, broiler performance, chlorine, water quality, 125*
Waterers per cage, cage shape, layer performance, nipple waterer, waterer location, 89*
Watering device, glomerular size, kidney anatomy, pullet strain, 31*
Wattle reaction, genetic analysis, *Staphylococcus aureus*, 70*
Weathering, chemical composition, layer diet, naked oats, 115*
Weight constant, age constant, body weight, feed efficiency, heritability, 1497
Weight correction, age and weight-constant efficiency, broiler growth, feed efficiency, feed intake, 565
Weight gain, abdominal fat, broiler sire strain, feed conversion, slaughter yield, 16
antibiotic, broiler performance, feed conversion, 70*
barley, chick performance, enzyme supplementation,

viscosity, 147*
beak trimming, behavior, feed consumption, pullet, 109*
broiler chick performance, management, semi-isolation brooding, 96*
broiler diet, energy use, feed efficiency, light source, 47*
broiler performance, carcass composition, isoproterenol, 95*
fasting, heat stress, potassium chloride, 36*
chick diet, copper, peanut skins, zinc, 51*
egg type pullet, mash, pellets *vs.* mash, temperature, 375
growth curve, multiphasic analysis, 33
Welfare, caged layer, despotism, dominance, peck order, 1008
Western blot analysis, antibody, immune response, layer, northern fowl mite, 549
Wet bulb temperature, broiler chick hatch weight, dry bulb temperature, hatching time, 163*
dry bulb temperature, hatch weight, 37*
Wet feed, dry feed, egg production, heat stress, 164*
Wheat, abdominal fat pad, broiler performance, perlite, 33*
Wheat contamination, body weight, chick growth, deoxynivalenol, organ weight, 1778†
Wheat middlings, dietary level, egg production, Haugh units, 1329
Wheat screenings, body weight, broiler, feed conversion, 615
White blood cell, antibody production, immune response, light regimen, thymidine uptake, 114*
White lupine, dehulling, meat flavor, turkey growth, 596
Whole body autoradiography, egg, ochratoxin A, quail-Japanese, tissue distribution, 707
Wing restraint, broiler breast meat, tenderness, textural quality, 275
Withdrawal period, abdominal fat pad, adipocyte size, broiler, monensin, 339†
Xanthophyll, Bermudagrass, egg production, egg weight, yolk color, 1154
broiler performance, broiler pigmentation, pigment source, 544
carotene, drying method, grass-clipping length, turf Bermudagrass, 1644†
egg yolk pigmentation, layer, 126*
Yellow grease, broiler chick diet, fat blends, poultry fat, 27*
Yield, composition, kamaboko-like products, mechanically deboned meat, poultry meat color, 112*
cooking method, marinated chicken fillet, moisture content, volatile analysis, 128*
Yolk, carotenoid composition, egg, lutein, serum, 608
Yolk color, Bermudagrass, egg production, egg weight, xanthophyll, 1154
Yolk membrane, lipase activity, pancreas, turkey poult, 1089
Yolk position, age of bird, egg quality, egg storage position, shell treatment, 75*
Yolk sac, glucose, glycogen, intestine, turkey poult, 493
Yolk stalk, broiler, intestine, location, Meckel's diverticulum, 677†
Youth development, Coturnix International, educational program, goals and strategy, 167*
Youth programs, FFA and 4-H, poultry majors, poultry programs of the future, student recruitment, 886
Z chromosome, avian leukosis virus, early feathering, endogenous virus, genes, 191
Zeolite, broiler chick, tibial dyschondroplasia, vitamin A, 113
calcium, chloride, phosphorus, tibial dyschondroplasia, 1436
egg shell quality, egg specific gravity, sodium aluminosilicate, 577
Zeolite-A, bone calcium, chick, coccidiosis, feed efficiency, 172*
broiler breeder egg, fertility, hatchability, 19*
broiler performance, chick weight, egg injection, 1*
chick weight gain, coccidiosis, duodenal pH, monensin, 171*
Zinc, aluminum, egg production, molt-induced, quail-Japanese, 1157
chick, copper, histidine, 47*
chick diet, copper, peanut skins, weight gain, 51*
duodenum, feed consumption, layer, molt-induced, 58*
egg weight, layer, molt-induced, 5*
force resting, iron, layer performance, 3*
rooster, spermatozoal integrity, 103*
Zinc deficiency, chick, immune response, 139*